

Cow herd vaccination programs, like calf vaccination programs<sup>2</sup>, are an important component of a herd health program. Vaccination protocols should be designed with specific management systems and production goals in mind. Cows are generally vaccinated for two reasons: 1) to prevent disease in the cow and 2) to pass immunity to the unborn calf.

## Preventing Disease in the Cow and Passing Immunity to Calves

Some cow vaccinations, like blackleg (clostridial), are given to protect the cow from diseases that affect her directly. Other vaccines are given to develop immunity in her unborn calf, which later protects it from diseases such as rotavirus, coronavirus, or *Escherichia coli* (*E. coli*), agents that cause scours in newborn calves. The need for specific vaccinations of this type may vary depending on the disease prevalence in a given area and the age of the calf when a disease will potentially occur.

A third category of vaccines affect the cow directly by preventing disease and affect the calf indirectly by preventing abortion. Examples of diseases these vaccines protect against are viral diseases like infectious bovine rhinotracheitis (IBR) and bovine viral diarrhea (BVD), bacterial diseases like vibriosis (bovine venereal campylobacteriosis) and leptospirosis, and protozoal diseases like neospora and trichomoniasis. In New Mexico, most pregnancy losses occur between mid-December and mid-February. Therefore, it may be preferable to vaccinate cows in the fall when cows are gathered for pregnancy testing so the immunity stimulated

by the vaccination will peak just prior to the greatest disease challenge to the pregnancy. In the fall, cows are generally in the best body condition of the production year, so under-nutrition is less likely to impact cows' response to vaccination.

## Vaccination Protocols

If cows are on a spring vaccination (fall calving) schedule, modifications to the fall vaccination program shown here may be needed. It is important to note that there is a wide variety of vaccine types, combinations, and uses. Labeled directions must be strictly followed because some vaccines have narrow parameters of efficacy. For example, trichomoniasis vaccine must be given just prior to the breeding season because the duration of immunity from vaccination is very short. If this vaccine is administered at a different time its efficacy may be compromised.

It is also important that bulls are included in the cow herd vaccination program because they can be a reservoir for some diseases. However, in the case of certain diseases, such as trichomoniasis, existing vaccines are not effective in bulls. This highlights the need to consult your veterinarian if you have questions about your vaccination program.

Vaccination is a tool that can be used very effectively to help reduce the incidence of disease in a cow herd. As with all tools, vaccinations must be administered correctly to get the full benefit. Proper timing, route of administration, dosage, and type of vaccine are paramount to preventing disease. Beef quality assurance guidelines should be followed whenever vaccines or medications are administered.

<sup>1</sup> Respectively, Extension Veterinarian, Extension Livestock Specialist, and Extension Associate, Department of Extension Animal Sciences and Natural Resources, New Mexico State University, Las Cruces.

<sup>2</sup> For information about calf vaccination programs, see New Mexico State University Cooperative Extension Guide B-223, *Calf Vaccination Guidelines*, available at [http://cahe.nmsu.edu/pubs/\\_b/B-223.pdf](http://cahe.nmsu.edu/pubs/_b/B-223.pdf).

Your veterinarian can help you develop a vaccination program specifically for your operation and geographic area. The following protocols are suggested for a fall vaccination program:

### **Cows and Bulls**

- Long-acting viral vaccine that includes at least IBR and BVD, and may include parainfluenza-3 virus (PI3) and bovine respiratory syncytial virus (BRSV)
- Long-acting campylobacter fetus (vibrio)/leptospirosis (lepto) vaccine
- Dewormer
- 7-way clostridial booster (optional)
- Other vaccines if necessary in your area

**Note:** In the fall, bred replacement heifers should be vaccinated using the same protocol as the mature cows.

### **Replacement Heifers (pre-breeding)**

- Modified-live IBR, BVD, PI3, BRSV (initial at 2 to 3 months of age, plus booster around weaning)
- Vibrio and 5-way leptospirosis (initial at 2 to 3 months of age, plus booster around weaning)
- Dewormer
- 7-way clostridial booster (optional)
- Brucellosis (optional; if administered it must be given by an accredited veterinarian and prior to 1 year of age)
- Other vaccines if necessary for your area

Contents of publications may be freely reproduced for educational purposes. All other rights reserved. For permission to use publications for other purposes, contact [pubs@nmsu.edu](mailto:pubs@nmsu.edu) or the authors listed on the publication.

New Mexico State University is an equal opportunity/affirmative action employer and educator. NMSU and the U.S. Department of Agriculture cooperating.