

Department of
Plant and Environmental Sciences

ABSTRACT

Few options of warm-season annual legumes are available for the higher elevations of semiarid, subtropical regions such as the Southern High Plains of the USA; therefore, candidate species must be explored for these environments. Pigeonpea [*Cajanus cajan* (L.) Millspaugh] has not been evaluated for forage in this environment, and pigeonpea forage yield and nutritive value have not been compared with those of cowpea [*Vigna unguiculata* (L.) Walpers], which is currently the most commonly used warm-season annual legume in the region. Pigeonpea (GA-2) and cowpea (Iron and Clay) were compared in separately analyzed randomized complete block studies conducted in 2006 and 2007 at New Mexico State University's Agricultural Science Center at Tatum using different fields and irrigation techniques each year. Cowpea was superior to pigeonpea for forage yield in 2006 and had numerically higher yield in 2007 [6.39 vs. 3.59 Mg ha⁻¹ for cowpea and pigeonpea, respectively, in 2006 (P <0.01) and 5.07 vs. 4.06 Mg ha⁻¹ in 2007 (P <0.30)]. Forage nutritive value of cowpea was greater than pigeonpea in both years as estimated by crude protein [189 vs. 156 g kg⁻¹ in 2006 (P <0.05) and 130 vs. 88 g kg⁻¹ in 2007 (P <0.01)] and neutral detergent fiber [323 vs. 467 g kg⁻¹ in 2006 (P <0.05) and 440 vs. 505 g kg⁻¹ in 2007 (P <0.05)]. Based on these results, cowpea remains the better option than pigeonpea for forage production systems under irrigation in the Southern

High Plains of the USA and similar higher elevation, semiarid, subtropical environments.

INTRODUCTION

Forage sorghum [*Sorghum bicolor* (L.) Moench] and sorghum x sudangrass (*S. bicolor* var. *Sudanese*) are important warm-season annual forage grasses in the higher elevations of the semiarid, subtropical Southern High Plains of the USA (eastern New Mexico and Texas Panhandle) because they are relatively drought-tolerant and productive. However, there are limitations regarding crude protein (CP) concentration and other nutritive value components, such as fiber and digestibility, and livestock may require supplementation if these components are too low (Mislevy et al., 2005; Contreras-Govea et al., 2009). Some of these deficiencies in nutritive value can be alleviated by including legumes in the forage system (Muir, 2002; Lauriault and Kirksey, 2004; Contreras-Govea et al., 2009). Few warm-season annual legumes are adapted to humid, subtropical regions (Mislevy et al., 2005). There are even fewer options for semiarid, subtropical regions, and additional potential forage legumes must be explored (Muir, 2002; Lauriault and Kirksey, 2004; Contreras-Govea et al., 2009).

Soybean (*Glycine max* L.) is widely adapted; however, drought limits its forage yields (Ishibashi et al., 2003; Mislevy et al., 2005), and the combination of high temperature and low humidity in the Southern High Plains increases the irrigation requirement for soybean and reduces grain yield (C. Trostle, personal communication, 2009).

¹Respectively, Forage Agronomist, Agricultural Science Center at Tatum (corresponding author; lmlaur@nmsu.edu); Crop Stress Physiologist, Agricultural Science Center at Clovis; Extension Agronomist, Agricultural Science Center at Clovis; and Agricultural Biometrician, Agricultural Biometrics Service, Agricultural Experiment Station, all of New Mexico State University.

Cowpea [*Vigna unguiculata* (L.) Walpers] is well adapted to low soil moisture situations (Muir, 2002), but not wet soils (Mislevy et al., 2005). Lauriault and Kirksey (2007) found that black-eyed cowpea grown for edible dry beans was well adapted to the Southern High Plains, with no grain yield reduction if planted before mid-June or if furrow irrigation or precipitation were timely, albeit limited. Data published by Contreras-Govea et al. (2009) indicate that forage cowpea has the potential to significantly improve nutritive value of monoculture sorghum forages.

Pigeonpea [*Cajanus cajan* (L.) Millspaugh] is also a widely adapted legume, but it prefers climates with an annual average temperature of 19°C and soils with 240 mm available water holding capacity at higher elevations (Mligo and Craufurd, 2005), which are prevalent in the Southern High Plains. Rao et al. (2003) and Mligo and Craufurd (2005) found that pigeonpea was least adapted to wet soil conditions, similar to cowpea (Mislevy et al., 2005). Consequently, pigeonpea has recently been promoted for grain and forage in the southern Great Plains (Rao et al., 2003; Bidlack et al., 2006; Rao and Dao, 2008). But that region has lower elevations than the Southern High Plains [approximately 400 m (Muir, 2002; Rao et al., 2003; Bidlack et al., 2006) vs. approximately 1,300 m (Lauriault and Kirksey, 2004; Contreras-Govea et al., 2009) for the southern Great Plains and the Southern High Plains, respectively] and higher precipitation [May through September, 500 mm in the southern Great Plains (Rao et al., 2003) vs. 280 mm in the Southern High Plains (Kirksey et al., 2003)].

No data are available about pigeonpea performance in the Southern High Plains of the USA. Consequently, the objective of this study was to compare forage yield and nutritive value of pigeonpea and cowpea in the Southern High Plains environment.

MATERIALS AND METHODS

Studies were conducted in 2006 and 2007 at the New Mexico State University Agricultural Science Center at Tucumcari, NM (35.20° N, 103.68° W; elev. 1,247 m), to compare forage production and nutritive value of 'GA-2' pigeonpea (University

of Georgia, Athens, GA) to that of 'Iron & Clay' cowpea (Turner Seed, Breckinridge, TX). Different fields with different soil types and irrigation techniques were used each year. The change in fields was warranted to avoid volunteerism by the crops from the previous year and a change in available irrigation techniques. Seeding rates were calculated to establish 150,000 and 111,000 plants ha⁻¹ for cowpea (Mislevy et al., 2005) and pigeonpea (Bidlack et al., 2006), respectively. Uninoculated seed was used for both studies because it was speculated that nodulation by either species could be poor locally due to soil factors. The average annual temperature and precipitation for this location are 14.4°C and 404.1 mm, respectively (Kirksey et al., 2003). The precipitation pattern in the area is continental, with approximately 80% (302.1 mm on average) falling from April through September (Kirksey et al., 2003). Weather data were collected from a National Weather Service station located within 1 km of the study areas (Figure 1).

In 2006, the soil was Caney fine sandy loam (fine-loamy, mixed, thermic Ustollic Haplargid), and the previous crop was sorghum x sudangrass that had followed a 7-year-old alfalfa (*Medicago sativa* L.) -grass mixture. Soil testing indicated a pH of 8.4 and N, P, and K levels of 10, 19, and 210 ppm, respectively, in the surface 30 cm. The seedbed was conventionally tilled and formed into beds on 0.91-m centers for furrow irrigation. Fertilizer (116-58-0 kg N-P₂O₅-K₂O ha⁻¹) was applied preplant and incorporated, and was based on soil test recommendations for sorghum forage and because the legume seed was not inoculated. There were four randomized complete blocks in which individual plots were 15.24 x 3.66 m, with a single row planted down the center of each of the 4 beds. Planting took place 25 May using a 4-row John Deere flex-planter with a seed-metering cone on each planting unit. Furrow irrigations were applied every other week from planting until mid-August to achieve field capacity with each application for a total estimated application of approximately 457 mm. Micronutrient deficiency was apparent early in the growing season, and was most pronounced on cowpea. Consequently, a commercially available micronutrient solution (HiYield Liquid Iron and Other Micronutrients, Voluntary Purchasing Groups, Bonham, TX) containing Cu, Fe, Mn, S, and Zn




Figure 1. Mean monthly air temperatures and total monthly precipitation during 2006 and 2007 and the long-term (1905–2005) averages at Tucumcari, NM, in the semiarid, subtropical Southern High Plains of the USA.

was applied foliarly to all plots at the labeled rate after two of the irrigations.

In 2007, the soil type was Redona fine sandy loam (fine-loamy, mixed, superactive, thermic Ustic Calcicargid) in a field that had been in native short prairie grasses for several decades. The pH of this soil was 7.2 and N, P, and K levels were 1, 2, and 19 ppm, respectively. There were three randomized complete blocks. Individual plots were 7.62 x 3.05 m with 4 rows spaced 0.76 m apart. Cowpea and pigeonpea were planted into a conventionally tilled flat seedbed under an overhead irrigation system on 30 May using the same planter that had been used in 2006, adjusted to the 0.76-m row spacing. Fertilizer (201-90-67 kg N-P₂O₅-K₂O ha⁻¹) was applied preplant and incorporated based on soil test recommendations for sorghum forage. Sprinkler irrigations (12.7 mm) were applied approximately every 3 d until harvest, resulting in approximately 470 mm applied

for the growing season. Micronutrient deficiency was not observed.

A section (3.05 m in 2006 and 2.0 m in 2007) of a center row of each plot was harvested with hand pruners, leaving a 5-cm stubble on 25 September 2006 and on 14 or 19 September 2007 for pigeonpea and cowpea, respectively, when the pods from the first flush of blooms began turning brown, which was slightly later than the harvest timing described by Mislavy et al. (2005), but within the range of harvest dates used by Rao et al. (2003). In 2006, harvested material from each plot was chopped using a stationary Gehl Model FH188 harvester; a lawn chipper/shredder was used in 2007. Plot weights were measured in the field. A subsample of chopped material (approximately 300 g) from each plot was immediately weighed, dried in a forced-air oven at 65°C for 48 h, and reweighed to determine dry matter (DM) concentration and to convert field

Table 1. Dry Matter Yield and Nutritive Value of Cowpea and Pigeonpea and Results of Statistical Analyses By Year¹

Variable	Units	Species		P-value
		Cowpea	Pigeonpea	
2006				
DM ² yield	Mg DM ha ⁻¹	6.39	3.59	0.0034
CP	g kg ⁻¹	189	130	0.0361
ADF	g kg ⁻¹	325	386	0.0675
NDF	g kg ⁻¹	329	467	0.0125
TDN	g kg ⁻¹	655	586	0.0686
2007				
DM yield	Mg DM ha ⁻¹	5.07	4.06	0.2959
CP	g kg ⁻¹	156	88	0.0046
ADF	g kg ⁻¹	402	402	1.0000
NDF	g kg ⁻¹	440	505	0.0122
TDN	g kg ⁻¹	567	568	0.9866

¹The study was conducted in the semiarid, subtropical Southern High Plains at Tucumcari, NM, USA, in different fields in 2006 and 2007.

²DM, CP, ADF, NDF, and TDN signify dry matter, crude protein, acid detergent fiber, neutral detergent fiber, and total digestible nutrients, respectively. Data are the means of four randomized complete blocks in 2006 and three randomized complete blocks in 2007.

weights to yield as Mg DM ha⁻¹. Dried samples were ground to pass a 1-mm screen and submitted to Ward Laboratories (Kearney, NE) for estimation of forage nutritive value [CP, acid detergent fiber (ADF), and neutral detergent fiber (NDF)] by near infrared spectroscopy (NIRS) using a universal equation developed from a broad spectrum of annual and perennial forage species. Total digestible nutrients (TDN) was calculated as part of the NIRS analysis.

Forage DM yield and nutritive value data were analyzed by year using SAS PROC MIXED procedures (SAS Institute, 2007) to determine if differences existed ($P < 0.05$) between species. Replicate was considered random.

RESULTS AND DISCUSSION

Dry matter yield. Cowpea outyielded pigeonpea in 2006 and had numerically higher yield in 2007 (Table 1). Forage yields in this study compared well with those measured by others. In the southern Great Plains (500 mm average May through September precipitation), Bidlack et al. (2006) reported rainfed pigeonpea yields of 1.9 to 4.6 Mg DM ha⁻¹, with a mean yield over three years of 2.3 Mg DM ha⁻¹ at populations and growing season length equivalent to the present study. In the humid subtropical southeastern USA, Mislevy et al. (2005) reported cowpea yields of 6.5 Mg DM ha⁻¹, while at the same location, Rao et al. (2003) measured pigeonpea leaf and stem yield of 6.0 Mg DM ha⁻¹. To the south of Rao et

al. (2003) and Bidlack et al. (2006), Muir (2002) measured irrigated hand-plucked forage cowpea yields of 4.3 to 5.6 Mg DM ha⁻¹. While pigeonpea was not included in that study, 290 mm of April to September precipitation was not sufficient for adequate forage production of several annual legumes, including cowpea (Muir, 2002). April through September precipitation in 2006 and 2007 at Tucumcari was 343 and 177 mm, respectively, in addition to the irrigation applications. With the application of irrigation water each year, available soil water should have been sufficient (>240 mm) to optimize pigeonpea yield (Mligo and Craufurd, 2005). Factors associated with lower productivity by pigeonpea compared to studies conducted elsewhere could include climate and soil differences. Annual temperatures at Tucumcari are approximately 5°C lower than the location of Rao et al. (2003) and the higher elevations used by Mligo and Craufurd (2005). Additionally, the soils used by those researchers (Rao et al., 2003; Mligo and Craufurd, 2005) were nearly neutral to heavily acidic (ISRIC, 2001), while the soils in the present study were nearly neutral to alkaline, with pH of 7.2 (2007) and 8.4 (2006).

Forage nutritive value. Cowpea nutritive value was higher in this study as indicated by greater CP and lower NDF than pigeonpea (Table 1). Nutritive value data estimated by NIRS for the present study are also consistent with those reported in the literature. Mislevy et al. (2005) measured cowpea CP of 140 to 220 g kg⁻¹ in a one-cut system.

For pigeonpea forage, Karachi and Matata (1996) reported CP concentrations of 205 g kg⁻¹, which is considerably higher than pigeonpea CP in the present study, while Rao et al. (2003) measured 135 g kg⁻¹ CP and 580 g kg⁻¹ digestibility, which are consistent with the data reported in Table 1 for CP. Still, the direct comparison of cowpea and pigeonpea in the present study demonstrates higher CP and lower NDF for cowpea than for pigeonpea. Lower NDF has been associated with greater intake potential by livestock (Chapko et al., 1991). Chapko et al. (1991) concluded that oat forage was higher in nutritive value than barley forage based on a 17 g kg⁻¹ difference in CP and a 67 g kg⁻¹ difference in NDF, both of which are considerably lower than the differences measured between cowpea and pigeonpea for those same variables in the present study (Table 1).

CONCLUSION

These data indicate that cowpea forage yield and nutritive value are superior to pigeonpea under irrigation in the Southern High Plains of the USA and similar higher elevation, semiarid, subtropical environments. Lower productivity of pigeonpea in this environment compared to other locations described in the literature may have been related to differences in climatic and soil effects.

ACKNOWLEDGMENTS

We gratefully acknowledge the provision of pigeonpea seed by S.C. Pathak, Department of Horticulture, University of Georgia, Tifton; technical and field assistance of Rusty Brake, Tamie Elam, Calvin Henson, Martin Mead, and Larry Perkins; secretarial assistance of Patty Cooksey; and the staff with the NMSU Library Document Delivery Service. Salaries and research support were provided by state and federal funds appropriated to the New Mexico State University Agricultural Experiment Station.

LITERATURE CITED

- Bidlack, J.E., A. Middick, D. Shantz, C.T. MacKown, R.D. Williams, and S.C. Rao. 2006. Weed control in a pigeon pea–wheat cropping system. *Field Crops Research*, 96, 63-70.
- Chapko, L.B., M.A. Brinkman, and K.A. Albrecht. 1991. Oat, oat–pea, barley, and barley–pea for forage yield, forage quality, and alfalfa establishment. *Journal of Production Agriculture*, 4, 486-491.
- Contreras-Govea, F.E., L.M. Lauriault, M.A. Marsalis, S. Angadi, and N. Puppala. 2009. Performance of forage sorghum–legume mixtures in Southern High Plains, USA [Online]. *Forage and Grazinglands*. DOI: 10.1094/FG-2009-0401-01-RS.
- Ishibashi, T., C.H. Sneller, and G. Shannon. 2003. Soybean yield potential and phenology in the ultra-short-season production system. *Agronomy Journal*, 95, 1,082-1,087.
- ISRIC. 2001. Major soils of the world [CD-ROM]. International Soil Reference and Information Centre. Wageningen University, The Netherlands.
- Karachi, M., and Z. Matata. 1996. Preliminary evaluation of pigeon-pea [*Cajanus cajan* (L.) Millsp.] accessions for forage production and nutritive value and seed yield in western Tanzania. *Tropical Agriculture*, 73, 253-258.
- Kirksey, R.E., L.M. Lauriault, and P.L. Cooksey. 2003. *Weather observations at the Agricultural Science Center at Tucumcari, 1905-2002* [Research Report 751]. Las Cruces: New Mexico State University Agricultural Experiment Station.
- Lauriault, L.M., and R.E. Kirksey. 2007. *Planting date and furrow-irrigation effects on cowpea for edible dry beans, Southern High Plains, USA* [Research Report 757]. Las Cruces: New Mexico State University Agricultural Experiment Station.
- Lauriault, L.M., and R.E. Kirksey. 2004. Yield and nutritive value of irrigated winter cereal forage grass-legume intercrops in the Southern High Plains, USA. *Agronomy Journal*, 96, 352-358.
- Mislevy, P., A.R. Blount, F.G. Martin, and B.T. Scully. 2005. Soybean and clay cowpea grown for forage production in the subtropics [Online]. *Crop Management*. DOI: 10.1094/CM-2005-0926-01-RS.
- Mligo, J.K., and P.Q. Craufurd. 2005. Adaptation and yield of pigeonpea in different environments in Tanzania. *Field Crops Research*, 94, 43-53.
- Muir, J.P. 2002. Hand-plucked forage yield and quality and seed production from annual and short-lived perennial warm-season legumes fertilized with composted manure. *Crop Science*, 42, 897-904.
- Rao, S.C., and T.H. Dao. 2008. Relationships between immobilized phosphorus uptake in two grain legumes and soil bioactive phosphorus pools in fertilized and manure-amended soil. *Agronomy Journal*, 100, 1,535-1,540.

Rao, S.C., W.A. Phillips, H.S. Mayeux, and S.C. Phatak. 2003. Potential grain and forage production of early maturing pigeonpea in the southern Great Plains. *Crop Science*, 43, 2,212-2,217

SAS Institute. 2007. The SAS system for Windows, release 9.1 [Computer software]. SAS Institute, Inc., Cary, NC.



***Leonard Lauriault** is a Forage Agronomist at New Mexico State University. He earned his M.S. at the University of Kentucky. Leonard's research program focuses on management and utilization of irrigated forage crops, including alfalfa, for pasture, hay, and biofuels.*

Brand names appearing in publications are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. Persons using such products assume responsibility for their use in accordance with current label directions of the manufacturer.

Contents of publications may be freely reproduced for educational purposes. All other rights reserved. For permission to use publications for other purposes, contact pubs@nmsu.edu or the authors listed on the last page of the publication.

New Mexico State University is an equal opportunity/affirmative action employer and educator. NMSU and the U.S. Department of Agriculture cooperating.