



The 2015 New Mexico Alfalfa Variety Test Report



Agricultural Experiment Station
College of Agricultural, Consumer
and Environmental Sciences



The 2015 New Mexico Alfalfa Variety Test Report

Leonard Lauriault, Ian Ray, Chris Pierce, Owen Burney, Robert Flynn, Mark Marsalis, Mick O'Neill, and Margaret West¹

Introduction

In 2015, 220,000 acres of alfalfa (*Medicago sativa*) were in production in New Mexico, which was the same as in 2014. Hay yields were estimated at 1,008,000 tons. At a January through November 2015 average of \$259/ton (up slightly from 2014), estimated gross returns from just over 1 million tons of alfalfa hay produced in 2015 will total nearly \$264 million. This is below the \$293 million received in 2014, but alfalfa hay continues as New Mexico's No. 1 cash crop (New Mexico Agricultural Statistics Service, www.nass.usda.gov/nm). Alfalfa also is the legume of choice in irrigated perennial pastures. Whether used as pasture or hay, the value of alfalfa to New Mexico is greatly magnified by its contribution to livestock production and receipts from the sale of meat, milk, and other products generated by livestock enterprises.

Choosing a good alfalfa variety is a key step in establishing a highly productive stand of alfalfa, whether for hay or pasture. Differences between the highest- and lowest-yielding varieties in irrigated tests included in this report ranged from 0.44 to 2.17 tons per acre in 2015. If sold as hay, this translates to a difference in returns of \$113 to \$562 per acre due to variety, or an increase of at least \$25.1 million for the industry in 2015 alone.

This report, which is a collaborative effort of New Mexico State University scientists at agricultural science centers throughout the state, provides yield data for alfalfa varieties included in yield trials in New Mexico. While consistently high yields compared to other varieties over a number of years and locations within a region is the best indication of varietal adaptation and persistence, other factors should be considered in the variety selection process (see NMSU's Cooperative Extension Service Circular 654, *Selecting alfalfa varieties for New Mexico*). In addition to fall dormancy and winter hardiness, high levels of pest resistance are critical to protecting an alfalfa stand for long-term production. Alfalfa grown in New Mexico should have at least a resistant (R) rating for bacterial wilt, Fusarium wilt, anthracnose, Phytophthora root rot, spotted alfalfa aphid, blue alfalfa aphid, pea aphid, stem nematode, and southern rootknot nematode. Seed quality also should be high. Selecting an alfalfa variety based on seed cost is a gamble producers often lose. To be assured of achieving a long-lasting, highly productive stand, buy either certified or Plant Variety Protected (PVP) seed, which guarantees the genetics and performance. The best choice of seed of any variety is one that was treated with a fungicide and nitrogen-fixing bacteria before it was bagged.

Description of Tests

Replicated alfalfa variety tests included in this report were conducted under research controls at NMSU's Agricultural Science Centers at Las Cruces (2014 normal, drought, and early termination irrigation studies), Artesia [2013, and 2014 (late spring planted)], Tucumcari (standard and Roundup Ready[®], both sown in 2012 irrigated with treated municipal wastewater), Los Lunas (2013), Mora (2013), and Farmington (2014).

Weather data for 2015 and the long-term averages from all locations are presented in table 1.

Yield data (on a dry matter basis) are presented in tables 2-11. Varieties are listed in order from highest to lowest average annual production. Yields are given by cutting for 2015 and by year for each production year. Statistical analyses were performed on all alfalfa yield data (including experimental entries) to determine if the apparent differences are truly due to variety or just to chance. The variety with the highest numerical yield in each column is marked with two asterisks (**), and those varieties not significantly different from that variety are marked with one asterisk (*). Those are the varieties from which to make an initial selection. Otherwise, to determine if two varieties are truly different, compare the difference between the two varieties to the Least Significant Difference (LSD) at the bottom of the column. If the difference is equal to or greater than the LSD, the varieties are truly different in yield when grown under the conditions at a given location. If NS is given for the LSD, there was no statistical difference between the highest and lowest yielding varieties. The Coefficient of Variation (CV), which is a measure of the variability of the data, is included for each column of means. Low variability (<20 percent) is desirable, and increased variability within a study results in higher CVs and larger LSDs. There might be a difference between previously published data and the data given in this publication for the same tests because of differences in the programs used for statistical analysis.

Table 12 summarizes information about proprietors, Roundup Ready genetics, fall dormancy, winter survival (measured in the northern United States), pest resistance, and yield performance across years and locations for all varieties currently included in NMSU's alfalfa variety testing program. Varieties are listed alphabetically by fall dormancy category. As in the data tables, the variety with the highest numerical yield in each column is marked with two asterisks (**), and those varieties not significantly different from that variety are marked with one asterisk (*). Remember good performance across several years and locations is the best indicator of broad adaptation, pest resistance, and persistence.

Seed labeled "common," "variety not stated," or "variety unknown", particularly that from other states, is of unknown genetic background and may or may not have the necessary disease or insect resistance. New Mexico Common and African Common seed used in all tests throughout the state has come from the same supplier and seed fields in New Mexico. Seed purchased from other dealers may or may not be of the same quality and performance.

Summary

Consistent production of high alfalfa yields is the result of selecting good varieties and implementing good management techniques. Soil fertility should be maintained at recommended levels based on soil tests, irrigation should be properly applied, weeds and insects should be controlled using appropriate cultural and/or chemical methods, and harvest management should allow sufficient time to restock root energy prior to winter.

¹Forage Crop Management Scientist and Superintendent, NMSU Agricultural Science Center at Tucumcari; Alfalfa Breeder, NMSU, Las Cruces; Forage Research Scientist, NMSU, Las Cruces; Superintendent, NMSU John T. Harrington Forestry Research Center at Mora; Extension Agronomist, NMSU Agricultural Science Center at Artesia; Extension Forage Specialist and Superintendent, NMSU Agricultural Science Center at Los Lunas; Agronomist, NMSU Agricultural Science Center at Farmington; and Agricultural Research Scientist, NMSU Agricultural Science Center at Farmington, respectively.

For dormant (FD 1 to 3) and semidormant (FD 4 to 6) varieties, a 6-week rest period before a dormancy-inducing freeze (27°F) is recommended to allow plants to replenish root reserves for winter survival and initiate spring growth, after which harvesting might be done either mechanically or by grazing. Non-dormant (FD 7 to 9) varieties also might benefit from this rest period. Removing fall growth is beneficial to reducing weevil populations the following year as eggs are laid in and overwinter in stems. Harvesting established stands at early bloom would result in 3 to 5 cuttings per year before initiation of the rest period in most areas of New Mexico. More dormant varieties might not produce yields that can be baled during the rest period; however, these can still be grazed. For further information about alfalfa management, refer to the other NMSU Agricultural Experiment Station and Cooperative Extension Service publications listed in table 13.

Acknowledgements

The authors express appreciation to the following for their significant contribution to The New Mexico Alfalfa Variety Testing Program by helping with planting, maintaining, harvesting, or other data collection, and data entry: Dallen Begay, Nathan Begay, Jason Box, Servando Bustillos, Patty Cooksey, Josh Foster, Charles Havlik, Jared Jennings, Shane Jennings, Dusty Lopez, Curtis Owen, Ruben Pacheco, Tom Place, Mario Segura, and the staff at University Communications who make publications such as this possible. Salaries and research support were provided by state and federal funds appropriated to the New Mexico Agricultural Experiment Station. Federal funds included those appropriated through the Hatch Act of 1887.

Table 1. Temperature and precipitation data for 2015 and the long-term averages for the New Mexico Alfalfa Variety Test locations.

Location Elevation Latitude	Las Cruces 3832 ft. 32° 12' N				Artesia 3366 ft. 32° 45' N				Tucumcari 4091 ft. 35° 12' N				Los Lunas 4840 ft. 34° 46' N				Mora 7303 ft. 35° 58' N				Farmington 5640 ft. 36° 41' N			
	Temp. (°F)		Precip. (in.)		Temp. (°F)		Precip. (in.)		Temp. (°F)		Precip. (in.)		Temp. (°F)		Precip. (in.)		Temp. (°F)		Precip. (in.)		Temp. (°F)		Precip. (in.)	
Month	2015	Avg.	2015	Avg.	2015	Avg.	2015	Avg.	2015	Avg.	2015	Avg.	2015	Avg.	2015	Avg.	2015	Avg.	2015	Avg.	2015	Avg.	2015	Avg.
Nov-14	48	49	0.68	0.53	46	49	1.63	0.53	45	47	0.37	0.71	43	43	0.34	0.46	38	42	0.64	0.21	41.7	41	0.31	0.64
Dec-14	45	41	0.13	0.68	43	41	0.22	0.51	40	39	0.38	0.65	38	35	1.03	0.53	33	32	0.62	0.48	34.5	31	0.88	0.46
Jan-15	41	40	0.70	0.56	37	40	1.32	0.39	36	38	1.44	0.42	35	35	0.99	0.38	34	33	0.7	0.11	32.7	30	1.22	0.49
Feb-15	50	45	0.03	0.37	44	45	0.23	0.42	42	42	0.89	0.48	43	40	0.21	0.41	36	33	1.04	0.30	39.7	36	0.93	0.47
Mar-15	57	52	0.29	0.22	51	52	0.12	0.43	50	49	0.38	0.76	51	47	0.36	0.49	43	43	1.21	0.29	46.9	43	0.31	0.65
Apr-15	62	60	0.43	0.21	60	60	0.92	0.62	58	57	1.93	1.14	55	55	0.87	0.45	48	49	0.57	0.31	51.1	51	0.61	0.58
May-15	67	69	0.10	0.29	66	69	2.47	1.20	62	66	4.02	1.94	62	63	1.47	0.45	52	60	4.43	0.54	57.5	60	1.70	0.50
Jun-15	80	78	0.00	0.72	78	78	0.83	1.40	76	76	2.07	1.95	77	72	0.33	0.54	65	67	3.95	1.33	71.7	70	1.80	0.23
Jul-15	80	80	0.94	1.36	82	80	1.19	1.76	79	79	7.56	2.73	77	77	3.60	1.39	68	69	3.21	3.34	73.7	76	0.91	0.81
Aug-15	80	78	0.78	2.29	81	78	0.38	1.67	78	77	2.03	2.76	77	75	0.81	1.67	68	68	5.93	2.19	74.1	74	1.14	1.04
Sep-15	75	71	0.23	1.38	76	71	0.93	1.81	76	71	1.31	1.61	71	67	0.54	1.17	65	62	0.5	1.81	68.4	66	0.51	1.11
Oct-15	63	61	2.75	0.91	63	61	4.49	1.16	61	60	2.36	1.30	60	56	1.75	1.05	53	51	4.18	0.64	58.6	54	0.04	0.96
Annual	62	60	7.06	9.52	60	60	14.73	11.90	59	58	24.74	16.45	57	55	9.41	8.99	50	51	26.98	11.55	54	53	10.36	7.94

Table 2. Dry matter yields (tons/acre) of alfalfa varieties sown October 7, 2014, at NMSU's Leyendecker Plant Science Research Center at Las Cruces and flood-irrigated twice per cutting to apply 52 inches (normal irrigation)†.

Variety Name	2015 Harvests						2015 Total
	28-Apr	12-Jun	16-Jul	12-Aug	9-Sep	6-Nov	
AmeriStand 803T	2.49	2.07*	1.64**	1.47*	1.28**	1.02**	10.07**
NM14BMHS1	2.88**	2.15*	1.56*	1.35*	1.16	0.95*	9.93*
NM14BMC0	2.61*	2.20**	1.62*	1.40*	1.16	0.96*	9.85*
NM14MaiHS3	2.60*	2.07*	1.58*	1.42*	1.16	0.97*	9.85*
Cisco II	2.35	2.11*	1.54*	1.36*	1.14	0.88	9.42*
NM14BMHS3	2.59	2.04*	1.49*	1.34	1.13	0.98*	9.42*
NM14GTAF07235	2.27	1.93	1.53*	1.51**	1.22*	0.90	9.42*
FSG903	2.31	2.01*	1.59*	1.30	1.15	0.91	9.33*
NM14BMHR2	2.55	2.03*	1.41	1.22	1.04	0.92	9.15
NM14ALWLHQ	2.26	1.95	1.45*	1.25	1.09	0.92	9.06
4N900	2.14	1.87	1.51*	1.40*	1.15	0.91	9.00
WL 440HQ	2.35	2.02*	1.40	1.19	1.04	0.94	8.99
WL 656HQ	2.20	1.87	1.49*	1.44*	1.17*	0.94	8.98
AFX096043	2.19	1.98*	1.52*	1.27	1.12	0.80	8.88
Wilson	2.02	1.92	1.44	1.39*	1.10	0.84	8.88
NM14MLLS2	2.4	2.04*	1.37	1.18	1.08	0.88	8.85
AFX149092	2.2	1.77	1.45*	1.32	1.13	0.91	8.80
NM1407227	2.43	1.88	1.36	1.23	1.03	0.72	8.75
NM14BM1008251	2.28	1.81	1.38	1.21	1.12	0.86	8.71
Malone	1.98	1.69	1.47*	1.48*	1.23*	0.85	8.65
AFX148091	2.12	1.84	1.39	1.30	1.10	0.90	8.60
NuMex Bill Melton	2.18	1.84	1.40	1.23	1.05	0.90	8.53
Sandpiper	2.08	1.93	1.37	1.23	1.04	0.91	8.52
57Q53	2.02	1.76	1.22	1.15	0.95	0.84	7.90
Mean	2.31	1.95	1.46	1.32	1.12	0.90	9.06
LSD (0.05)	0.29	0.23	0.19	0.17	0.11	0.08	0.81
CV%	8.83	8.29	9.45	9.06	7.15	6.05	6.08

†Data were analyzed using analysis of covariance where check plots of NuMex Bill Melton were used as the covariate.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 3. Dry matter yields (tons/acre) of alfalfa varieties sown October 7, 2014, at NMSU's Leyendecker Plant Science Research Center at Las Cruces and flood-irrigated once per cutting to apply 27.56 inches (drought irrigation)†.

Variety Name	2015 Harvests						2015 Total
	30-Apr	6-Jun	7-Jul	13-Aug	10-Sep	28-Oct	
NM14BMHS3	2.52*	1.30**	0.86**	1.28**	0.75*	0.70*	7.44**
NM14BMHS1	2.53**	1.30*	0.81*	1.23*	0.75*	0.68*	7.29*
AmeriStand 803T	2.30*	1.20*	0.85*	1.16*	0.83**	0.72**	7.07*
NM14BM1008251	2.27*	1.17*	0.82*	1.24*	0.78*	0.65*	6.95*
NM1407227	2.43*	1.13*	0.77*	1.11*	0.77*	0.64	6.85*
AFX149092	2.12*	1.20*	0.84*	1.19*	0.80*	0.65*	6.80*
4N900	2.21*	1.12*	0.84*	1.15*	0.81*	0.64	6.78*
FSG903	2.07*	1.17*	0.82*	1.15*	0.79*	0.69*	6.69*
NM14MalHS3	2.16*	1.24*	0.81*	1.11*	0.70	0.69*	6.67*
Wilson	1.95	1.10*	0.81*	1.18*	0.77*	0.63	6.45*
NM14GTAF07235	1.83	1.10*	0.82*	1.22*	0.80*	0.66*	6.44*
NM14BMC0	2.39*	1.00*	0.62	1.07*	0.68	0.62	6.39*
NM14ALWLHQ	1.82	1.10*	0.85*	1.19*	0.73*	0.63	6.31*
NuMex Bill Melton	1.95	1.07*	0.76*	1.17*	0.72	0.62	6.27*
NM14BMHR2	2.19*	0.91*	0.66	1.03*	0.65	0.63	6.07
57Q53	1.89	0.97*	0.74*	1.09*	0.68	0.56	5.95
WL 656HQ	1.70	0.99*	0.78*	1.05*	0.73*	0.62	5.86
NM14MLLS2	1.91	0.88*	0.67	1.11*	0.67	0.60	5.83
Malone	1.48	0.96*	0.80*	1.18*	0.78*	0.58	5.77
Cisco II	1.92	0.93*	0.57	1.06*	0.63	0.59	5.72
AFX096043	1.60	0.90*	0.77*	1.11*	0.76*	0.55	5.69
Sandpiper	1.87	0.91*	0.63	1.01*	0.68	0.55	5.64
AFX148091	1.58	0.84*	0.67	1.07*	0.70	0.62	5.50
WL 440HQ	1.61	0.89*	0.64	1.01*	0.60	0.55	5.32
Mean	2.01	1.06	0.76	1.13	0.73	0.63	6.32
LSD (0.05)	0.50	NS	0.16	NS	0.10	0.08	1.29
CV%	17.60	20.50	14.54	10.33	9.84	8.72	13.93

†Data were analyzed using analysis of covariance where check plots of NuMex Bill Melton were used as the covariate.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 4. Dry matter yields (tons/acre) of alfalfa varieties sown October 7, 2014, at NMSU's Leyendecker Plant Science Research Center at Las Cruces and flood-irrigated twice per cutting from March 31 until July 1 to apply 39.57 inches, after which irrigation was terminated (early termination irrigation)†.

Variety Name	2015 Harvests				2015 Total
	29-Apr	12-Jun	17-Jul	12-Aug	
NM14BMHS1	2.78*	2.28**	1.70**	0.45*	7.15**
NM14BMC0	2.84**	2.26*	1.62*	0.34	7.08*
NM14BMHR2	2.82*	2.22*	1.59*	0.34	6.96*
NM14MaIHS3	2.73*	2.18*	1.61*	0.37	6.89*
NM14MLLS2	2.71*	2.18*	1.56	0.32	6.80*
NM14BMHS3	2.68*	2.21*	1.59*	0.34	6.79*
NM1407227	2.66*	2.22*	1.50	0.28	6.69*
NM14ALWLHQ	2.34	2.21*	1.69*	0.39*	6.65*
AmeriStand 803T	2.35	2.14*	1.63*	0.46*	6.63*
FSG903	2.43	2.11	1.66*	0.43*	6.61
NM14BM1008251	2.35	2.21*	1.60*	0.44*	6.60
4N900	2.35	2.05	1.67*	0.41*	6.48
AFX149092	2.29	2.00	1.67*	0.48**	6.47
AFX096043	2.39	2.09	1.63*	0.32	6.45
AFX148091	2.27	2.10	1.62*	0.42*	6.45
Cisco II	2.32	2.19*	1.58*	0.31	6.39
NuMex Bill Melton	2.34	2.02	1.55	0.39*	6.30
Sandpiper	2.40	2.06	1.53	0.30	6.27
WL 656HQ	2.25	1.88	1.65*	0.46*	6.23
NM14GTAF07235	2.17	1.96	1.54	0.50*	6.18
WL 440HQ	2.24	2.09	1.53	0.33	6.18
Wilson	2.17	2.00	1.53	0.44*	6.16
57Q53	2.30	2.00	1.47	0.27	6.02
Malone	1.91	1.80	1.51	0.47*	5.63
Mean	2.42	2.10	1.59	0.38	6.50
LSD (0.05)	0.21	0.16	0.12	0.12	0.49
CV%	6.19	5.58	5.51	22.49	5.66

†Data were analyzed using analysis of covariance where check plots of NuMex Bill Melton were used as the covariate.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 5. Dry matter yields (tons/acre) of sprinkler-irrigated alfalfa varieties sown September 23, 2013, at NMSU's Agricultural Science Center at Artesia†.

Variety Name	2014	2015 Harvests					2015	2-Yr
	Total	8-May	19-Jun	17-Jul	20-Aug	29-Sep	Total	Average
NM Common	8.50*	2.29*	2.28*	2.22*	1.58*	1.47*	9.83*	9.16**
Malone	8.00*	2.20*	2.39**	2.55**	1.64*	1.30*	10.08**	9.04*
Dona Ana	7.99*	2.14*	2.17*	2.31*	1.61*	1.54*	9.76*	8.88*
56S82	8.63**	2.24*	2.21*	1.79*	1.34	1.42*	9.00*	8.82*
WL 535HQ	8.12*	2.22*	2.07*	2.13*	1.65*	1.34*	9.41*	8.76*
African Common	8.20*	2.17*	1.85*	2.30*	1.59*	1.31*	9.22*	8.71*
Artesian Sunrise	8.54*	2.05*	2.15*	2.05*	1.27	1.35*	8.87*	8.71*
Wilson	8.02*	2.04*	1.92*	2.06*	1.67**	1.55**	9.24*	8.63*
Zia	7.94*	2.15*	2.19*	2.14*	1.41*	1.40*	9.29*	8.61*
58N57	7.64	2.29*	2.19*	2.10*	1.66*	1.30*	9.53*	8.58*
DG9212	7.93*	1.98*	2.39**	1.98*	1.31	1.29*	8.94*	8.43*
55Q27	7.54	2.16*	2.27*	2.28*	1.33	1.24*	9.27*	8.41*
55VR05	6.88	2.37**	1.50*	2.17*	1.20	1.20*	8.45*	7.66*
Mean	7.99	2.18	2.12	2.16	1.48	1.36	9.30	8.64
LSD (0.05)	0.82	NS	NS	NS	0.31	NS	NS	NS
CV%	7.17	9.76	18.14	14.23	14.39	16.32	7.86	8.47

†Data were analyzed using analysis of variance.

2014 Harvest dates: 20-May, 18-Jul, 20-Aug, and 2-Oct

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 6. Dry matter yields (tons/acre) of sprinkler-irrigated alfalfa varieties sown March 31, 2014, at NMSU's Agricultural Science Center at Artesia†.

Variety Name	2014 Total	2015 Harvests				2015 Total	2-Yr Average
		26-May	13-Jul	12-Aug	30-Sep		
56S82	4.15*	2.22**	2.55**	1.81*	1.79*	8.37**	6.26**
NM Common	4.36*	1.72*	2.21*	2.14**	2.06**	8.13*	6.24*
Malone	4.28*	1.95*	2.16*	2.12*	1.74*	7.97*	6.13*
Wilson	4.30*	1.95*	1.90*	2.00*	1.99*	7.84*	6.07*
DG9212	4.45*	1.93*	1.83*	2.00*	1.91*	7.66*	6.05*
NuMex Bill Melton	4.06*	2.18*	2.19*	1.90*	1.76*	8.03*	6.04*
SW 8357	4.09*	1.78*	2.29*	2.08*	1.86*	8.00*	6.04*
SW 8208	4.49**	1.90*	1.91*	1.96*	1.74*	7.50*	6.00*
58N57	4.42*	1.79*	2.06*	1.88*	1.86*	7.57*	6.00*
SW 8421S	3.82*	2.01*	2.17*	2.05*	1.86*	8.09*	5.96*
SW 7410	4.18*	1.66*	2.20*	2.11*	1.72*	7.69*	5.93*
Artesian Sunrise	4.16*	1.74*	2.18*	1.87*	1.77*	7.56*	5.86*
Zia	4.32*	1.43*	1.93*	2.13*	1.89*	7.37*	5.85*
WL 535HQ	4.08*	1.78*	1.99*	1.96*	1.78*	7.51*	5.79*
African Common	4.28*	1.75*	1.73*	1.99*	1.72*	7.18*	5.73*
Dona Ana	3.98*	1.92*	1.78*	1.79*	1.92*	7.41*	5.70*
55VR05	3.68*	1.53*	2.00*	1.79*	1.69*	7.00*	5.34*
Mean	4.18	1.84	2.06	1.97	1.83	7.70	5.94
LSD (0.05)	NS	NS	NS	NS	NS	NS	NS
CV%	9.91	19.60	16.77	12.06	10.59	9.50	10.80

†Data were analyzed using analysis of variance.

2014 Harvest dates: 24-Jun, 2-Sep, and 3-Oct.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 7. Dry matter yields (tons/acre) of alfalfa varieties sown September 26, 2012, at NMSU's Agricultural Science Center at Tucumcari and sprinkler-irrigated twice per week with treated municipal wastewater.

Variety Name	2013 Total	2014 Total	2015 Harvests						2015 Total	3-Yr Average
			14-May	13-Jun	14-Jul	19-Aug	17-Sep	10-Nov		
Mallard	3.26	5.62*	1.14*	1.47**	0.25	1.46**	0.47*	0.35	5.13*	4.74**
56S82	3.90**	5.82*	1.28*	1.44*	0.24	1.33*	0.51*	0.49**	5.29**	4.71*
Malone	3.90**	6.26*	0.83	1.39*	0.47**	1.46**	0.34	0.40*	4.88*	4.69*
Integra 8400	3.00	5.98*	1.16*	1.31*	0.14	1.34*	0.42*	0.36*	4.72	4.66*
NM Common	3.20	5.80*	0.92	1.25*	0.40*	1.28*	0.54*	0.46*	4.84*	4.64*
NuMex Bill Melton	3.45*	6.28**	1.05	1.34*	0.25	1.43*	0.30	0.39*	4.76	4.61*
WL 454HQ.RR	3.23	5.87*	1.07	1.28*	0.17	1.30*	0.35*	0.43*	4.59	4.57*
Roadrunner	3.52*	6.06*	1.33**	1.39*	0.24	1.37*	0.56*	0.19	5.07*	4.52*
55Q27	2.93	5.53*	1.01	1.17*	0.13	1.35*	0.68**	0.37*	4.70	4.43*
Wilson	2.95	4.98	1.01	1.21*	0.21	1.28*	0.50*	0.46*	4.66	4.39*
Bluejay HR	2.88	5.86*	1.20*	1.33*	0.15	1.35*	0.18	0.13	4.33	4.36*
6422Q	2.72	5.66*	0.93	1.37*	0.15	1.29*	0.34	0.27	4.33	4.29*
54VR03	2.52	5.00	1.03	1.18*	0.10	1.16*	0.38*	0.32	4.16	4.27*
African Common	3.18	5.92*	0.85	1.22*	0.32	1.45*	0.34	0.40*	4.59	4.27*
Meadowlark	2.67	5.73*	1.21*	1.21*	0.11	1.39*	0.18	0.21	4.30	4.22*
Dona Ana	3.28	4.87	0.83	1.25*	0.21	1.29*	0.46*	0.44*	4.49	4.22*
HybriForce-2400	2.27	5.04	1.30*	1.27*	0.14	1.27*	0.35*	0.21	4.54	4.15*
54QR04	3.12	5.57*	0.90	1.23*	0.11	1.27*	0.29	0.28	4.08	4.13*
Bluejay 2	2.76	4.66	1.13*	1.11*	0.10	1.38*	0.25	0.16	4.13	3.88*
Mean	3.09	5.60	1.06	1.28	0.20	1.34	0.39	0.33	4.61	4.41
LSD (0.05)	0.60	0.91	0.23	NS	0.15	NS	0.34	0.14	0.70	NS
CV%	13.69	11.40	15.60	17.31	53.57	11.61	61.64	29.19	10.78	10.80

†Data for 2013 and 2014 were detrended using nearest neighbor analysis, and analyzed using analysis of variance; 2015 data were analyzed using analysis of variance only.

2013 Harvest dates: 5-Jun, 25-Jun, 1-Aug, 23-Sep, and 6-Nov.

2014 Harvests: 21-May, 1-Jul, 21-Jul, 19-Aug, 16-Sep, and 31-Oct.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 8. Dry matter yields (tons/acre) of Roundup Ready® alfalfa varieties sown September 26, 2012, at NMSU's Agricultural Science Center at Tucumcari and sprinkler-irrigated twice per week with treated municipal wastewater†.

Variety Name	2013 Total	2014 Total	2015 Harvests						2015 Total	3-Yr Average
			14-May	13-Jun	14-Jul	19-Aug	17-Sep	10-Nov		
6829R	4.00**	8.78**	1.13	1.81*	0.93**	1.52	0.62*	0.47*	6.48*	6.42**
R65BD278	3.77*	8.66*	1.29*	1.95**	0.78*	1.74*	0.59*	0.46*	6.80**	6.41*
R58HG236	3.22	8.59*	1.20*	1.82*	0.81*	1.63*	0.72**	0.44*	6.61*	6.14*
R78T823	3.72*	8.24*	1.21*	1.85*	0.79*	1.44	0.70*	0.47*	6.45*	6.14*
R66BX312	3.81*	8.28*	1.23*	1.68	0.84*	1.35	0.60*	0.49*	6.19	6.09*
R66BX320	3.45*	8.60*	1.09	1.75*	0.80*	1.40	0.65*	0.53**	6.20	6.08*
R57K138	2.96	8.59*	1.37*	1.83*	0.81*	1.61*	0.50	0.45*	6.57*	6.04*
R57OK217	2.71	8.56*	1.28*	1.73	0.79*	1.73*	0.59*	0.43	6.55*	5.94*
R57A136	3.39*	8.04*	1.28*	1.69	0.72*	1.57	0.62*	0.34	6.21	5.88*
R57W213	2.77	8.50*	1.29*	1.69	0.69*	1.46	0.58*	0.50*	6.21	5.82
RR57K337	3.43*	8.12*	1.23*	1.59	0.77*	1.41	0.52	0.35	5.86	5.80
R66BX311	3.29	7.97*	1.08	1.74	0.86*	1.35	0.65*	0.48*	6.15	5.80
R57OK216	2.64	8.46*	1.38**	1.77*	0.81*	1.36	0.58*	0.37	6.27	5.79
54QR04	2.57	8.26*	1.38**	1.91*	0.75*	1.43	0.59*	0.39	6.43*	5.76
R65BD277	3.18	7.89*	1.08	1.71	0.79*	1.54	0.55	0.46*	6.12	5.73
R65BD279	2.94	7.90*	1.19*	1.66	0.77*	1.63*	0.45	0.49*	6.19	5.68
R86X214	2.24	8.01*	1.28*	1.77*	0.76*	1.82**	0.54	0.40	6.56*	5.60
54VR03	2.81	7.27	1.31*	1.56	0.69*	1.59	0.51	0.39	6.05	5.38
Mean	3.07	8.18	1.23	1.74	0.78	1.52	0.58	0.43	6.27	5.84
LSD (0.05)	0.71	1.06	0.20	0.21	NS	0.23	0.15	0.10	0.45	0.57
CV%	16.31	9.14	11.62	8.50	16.86	10.80	17.99	15.63	5.01	11.93

†Data were detrended using nearest neighbor analysis, and analyzed using analysis of variance.

2013 Harvest dates: 5-Jun, 25-Jun, 1-Aug, 23-Sep, and 6-Nov.

2014 Harvests: 21-May, 1-Jul, 21-Jul, 19-Aug, 16-Sep, and 31-Oct.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 9. Dry matter yields (tons/acre) of flood-irrigated alfalfa varieties sown September 30, 2013, at NMSU's Agricultural Science Center at Los Lunas†.

Variety Name	2014 Total	2015 Harvests					2015 Total	2-Yr Average
		17-Jun	23-Jul	21-Aug	12-Oct	12-Nov		
NuMex Bill Melton	8.33***	2.71*	1.87*	1.70*	1.73**	0.56*	8.57*	8.45***
WL 454HQ.RR	7.28*	2.81*	1.93*	1.79**	1.57*	0.49	8.59**	7.93**
Artesian Sunrise	7.65**	2.84*	1.87*	1.61*	1.42	0.19	7.93*	7.79*
Meadowlark	7.13	3.13**	1.89*	1.62*	1.26	0.09	7.98*	7.56*
Wilson	7.53*	2.40	1.79*	1.60*	1.44	0.34	7.57	7.55*
Transition 6.10RR	7.29*	2.38	1.63*	1.53*	1.70*	0.48	7.71	7.50*
Malone	7.19*	2.40	1.70*	1.51*	1.61*	0.52*	7.73	7.46*
Dona Ana	7.15	2.26	1.70*	1.49*	1.67*	0.63**	7.73	7.44
Cimarron VL600	7.10	2.76*	1.80*	1.55*	1.20	0.22	7.52	7.31
WL 440HQ	6.87	2.57	1.77*	1.70*	1.38	0.26	7.68	7.27
PGI 424	7.13	2.46	1.83*	1.57*	1.34	0.12	7.31	7.22
NM Common	7.13	2.26	1.60*	1.53*	1.34	0.44	7.16	7.14
Roadrunner	6.79	2.51	1.96**	1.76*	1.17	0.08	7.47	7.13
55Q27	6.84	2.48	1.73*	1.47*	1.50*	0.16	7.34	7.09
HybriForce-2400	7.23*	2.46	1.57*	1.42*	1.27	0.09	6.80	7.02
Mallard	6.44	2.51	1.88*	1.52*	1.39	0.14	7.44	6.94
PGI 557	6.57	2.33	1.84*	1.48*	1.44	0.13	7.21	6.89
Archer III	6.45	2.34	1.80*	1.66*	1.37	0.17	7.33	6.89
Bluejay HR	6.63	2.61	1.69*	1.43*	1.00	0.11	6.83	6.73
DG4210	6.27	2.29	1.93*	1.55*	1.30	0.10	7.16	6.72
55VR05	6.06	2.16	1.80*	1.40*	1.28	0.11	6.75	6.41
Bluejay 2	5.67	2.57	1.72*	1.40*	1.16	0.08	6.93	6.30
Mean	6.94	2.51	1.78	1.56	1.39	0.25	7.49	7.21
LSD (0.05)	0.50	0.44	NS	NS	0.24	0.12	0.78	0.48
CV%	5.06	12.52	10.98	12.94	12.12	32.67	7.38	6.61

†Data were detrended using nearest neighbor analysis, and analyzed using analysis of variance.

2014 Harvest dates: 28-May, 30-Jun, 18-Aug, and 4-Nov.

***Highest numerical value in the column; significantly higher than all other varieties.

Second highest numerical value in the column where * is shown.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 10. Dry matter yields (tons/acre) of sprinkler-irrigated alfalfa varieties sown August 28, 2013, at NMSU's John T. Harrington Forestry Research Center at Morat.

Variety Name	2014 Total	2015 Harvests			2015 Total	2-Yr Average
		22-Jun	14-Aug	5-Nov		
Dona Ana	4.06**	1.03*	1.09*	0.24**	2.36*	3.21**
SS120	3.77*	1.16*	1.11*	0.16	2.43*	3.10*
55VR05	3.49*	1.19*	1.29**	0.15	2.63*	3.06*
WL 354HQ	3.32*	1.32*	1.20*	0.14	2.65*	2.99*
DG4210	3.30*	1.29*	1.26*	0.11	2.65*	2.97*
WL 319HQ	3.16*	1.29*	1.26*	0.12	2.66*	2.91*
FG27C102	3.28*	1.31*	1.05*	0.13	2.48*	2.88*
Wilson	3.00*	1.24*	1.02*	0.19*	2.44*	2.72*
55Q27	2.72*	1.36*	1.17*	0.15	2.69*	2.70*
Ladak	2.45*	1.44**	1.21*	0.16	2.80**	2.62*
Spredor 5	2.77*	1.30*	1.06*	0.09	2.44*	2.60*
Ranger	2.65*	1.16*	1.17*	0.14	2.47*	2.56*
Mean	3.16	1.26	1.15	0.15	2.56	2.86
LSD (0.05)	NS	NS	NS	0.07	NS	NS
CV%	24.25	18.83	26.51	31.93	16.65	24.85

†Data were detrended using nearest neighbor analysis, and analyzed using analysis of variance.

2014 Harvest dates: 22-Jul, 2- Sep, and 14-Oct.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 11. Dry matter yields (tons/acre) of sprinkler-irrigated alfalfa varieties sown August 21, 2014, at NMSU's Agricultural Science Center at Farmington†.

Variety Name	2015 Harvests				2015 Total
	2-Jun	2-Jul	17-Aug	10-Sep	
Roadrunner	4.27**	2.29*	2.43*	1.78*	10.77**
Mallard	4.04*	2.22*	2.57*	1.67	10.50*
Raven	3.65*	2.25*	2.66**	1.92**	10.48*
Archer III	4.20*	2.17*	2.49*	1.62	10.48*
Lahonton	4.04*	2.26*	2.36	1.67	10.32*
Ranger	3.80*	2.00*	2.63*	1.81*	10.23*
Arrowhead II	3.56*	2.31**	2.61*	1.70	10.18*
PGI 424	4.09*	2.02*	2.37	1.57	10.04*
4S417	3.77*	2.02*	2.52*	1.73*	10.04*
WL 363HQ	3.47*	2.17*	2.54*	1.72	9.90*
NM Common	3.70*	2.15*	2.29	1.75*	9.90*
AFX103009	3.61*	2.15*	2.46*	1.65	9.87*
MagnaGraze II	3.78*	2.07*	2.31	1.72	9.87*
WL 354HQ	3.96*	2.09*	2.20	1.62	9.86*
Mountaineer 2.0	3.62*	2.14*	2.43*	1.62	9.81*
54VR03	3.98*	2.01*	2.18	1.47	9.64
Gunner	3.59*	2.01*	2.35	1.65	9.61
GrandStand	3.72*	1.96*	2.27	1.58	9.52
Dona Ana	3.57*	1.97*	2.28	1.71	9.52
Wilson	3.37*	1.86*	2.34	1.76*	9.33
Malone	3.15*	1.73*	2.35	1.65	8.87
Zia	3.34*	1.62*	2.48*	1.42	8.87
Mean	3.74	2.06	2.41	1.67	9.89
LSD (0.05)	NS	NS	0.29	0.20	1.06
CV%	12.79	14.55	8.53	8.58	7.55

†Data were detrended using nearest neighbor analysis, and analyzed using analysis of variance.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 12. Characteristics and performance of alfalfa varieties across years and tests in New Mexico.		Varietal Characteristics ¹											Las Cruces			Artesia				Tucumcari			Los Lunas		Mora		Farmington				
													2014 ²							2012			2013		2013		2014				
		Pest resistance											N ³	D	ET	2013	2014	Std	RR [®]	2013	2013	2014									
		Variety	Proprietor	RR	FD	WS	BW	FW	AN	PRR	SAA	PA	BAA	SN	RKN	15 ⁴	15	15	14	15	14	15	13	14	15	13	14	15	14	15	14
Arrowhead II	Alforex Seeds		2	2	HR	HR	HR	HR	n/r	R	n/r	HR	n/r																		*
FG 27C102	Forage Genetics Int.		2	1	HR	HR	HR	h	n/r	R	n/r	n/r	n/r																*	*	
Spreader 5	Nexgrow Alfalfa		2	1	HR	HR	HR	n/r	R	n/r	n/r	n/r	n/r																*	*	
Ladak	USDA		3	n/r	R	n/r	n/r	n/r	n/r	n/r	n/r	R	n/r															*	**		
MagnaGraze	Alforex Seeds		3	2	HR	HR	R	HR	R	n/r	MR	MR	LR																		
MagnaGraze II	Sharp Brothers		3	2	HR	HR	HR	n/r	R	n/r	n/r	R	n/r																	*	
Ranger	USDA/Univ. of Nebraska		3	n/r	R	n/r	n/r	n/r	R	n/r	n/r	R	n/r															*	*	*	
SS120	Seed Solutions		3	3	HR	R	R	R	R	R	R	n/r	n/r															*	*		
WL 319HQ	W-L Research		3	1	HR	HR	HR	HR	R	n/r	HR	n/r	n/r															*	*		
4S417	Mycogen Seeds		4	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r																	*	
54QR04	Pioneer HiBred Int'l	Y	4	n/r	HR	HR	HR	HR	n/r	n/r	n/r	n/r	n/r									*	*	*	*	*					
54VR03	Pioneer HiBred Int'l	Y	4	n/r	HR	HR	HR	HR	n/r	HR	n/r	n/r	n/r																		
6422Q	Nexgrow Alfalfa		4	1	HR	HR	HR	HR	n/r	R	n/r	R	n/r									*									
Bluejay 2	Blue River Hybrids		4	2	HR	HR	HR	HR	R	HR	n/r	n/r	n/r																		
Bluejay HR	Blue River Hybrids		4	2	HR	HR	HR	HR	R	HR	n/r	R	n/r									*									
DG4210	Crop Production Services		4	1	HR	HR	HR	HR	HR	R	n/r	R	n/r															*	*		
GrandStand	Crop Production Services		4	2	HR	HR	HR	HR	R	HR	n/r	MR	n/r																		
HybriForce Mesa	Alforex Seeds		4	2	HR	HR	HR	HR	n/r	R	n/r	HR	R																		
HybriForce-2400	Sharp Brothers		4	2	HR	HR	HR	HR	n/r	n/r	n/r	HR	R														*				
Integra 8400	Wilbur-Ellis Company/Integra		4	2	HR	HR	HR	HR	n/r	HR	n/r	R	n/r									*									
Meadowlark	Blue River Hybrids		4	2	HR	HR	HR	HR	R	HR	n/r	R	n/r									*						*			
PGI 424	Alforex Seeds		4	2	HR	HR	HR	HR	R	R	n/r	R	n/r																	*	
Raven	Blue River Hybrids		4	2	HR	R	R	R	R	HR	R	R	n/r																	*	
Roadrunner	Blue River Hybrids		4	2	HR	HR	HR	HR	HR	LR	n/r	R	n/r								*	*	*							**	
WL 354HQ	W-L Research		4	1	HR	HR	HR	HR	HR	HR	n/r	R	n/r															*	*	*	
55Q27	Pioneer HiBred Int'l		5	n/r	HR	HR	HR	HR	R	R	n/r	HR	n/r				*				*							*	*		
55VR05	Pioneer HiBred Int'l	Y	5	n/r	HR	HR	HR	HR	n/r	n/r	n/r	HR	n/r				*	*	*									*	*		

¹RR=Roundup Ready if "Y"; WS=Winter Survival (1=No injury, 6=Dead plants), FD=Fall Dormancy (1 & 2 Very dormant; 3 & 4 Dormant; 5 Moderately dormant; 6 & 7 Semi-dormant; 8 & 9 Non-dormant; 10 & 11 Very non-dormant), BW=Bacterial wilt, PRR=Phytophthora root rot, FW=Fusarium wilt, AN=Anthracnose, SAA=Spotted alfalfa aphid, PA=Pea aphid, BAA=Blue alfalfa aphid, SN=Stem nematode, RKN=Rootknot nematode (southern); (S=Susceptible, LR=Low resistance, MR=Moderate resistance, R=Resistant, HR=High resistance, n/r indicates either that the variety was not rated for that characteristic or no rating was available).

²Establishment year.

³N, D, ET, STD, and RR[®] signify normal irrigation, drought irrigation, early termination irrigation, standard performance evaluation, and Roundup Ready[®] entries only, respectively.

⁴Harvest year.

Shaded boxes indicate that the variety was not in the test.

***Significantly higher than all other values in the column.

**Highest yielding variety in the test for that year, except for Los Lunas 2013 test when it is the second highest yielding variety.

*Not significantly different from the highest yielding variety in the test for that year, except for Los Lunas 2013 test in 2014 when it is not significantly different from the second highest yielding variety.

L.M. Lauriault, I.M. Ray, C.A. Pierce, O. Burney, R.P. Flynn, M.A. Marsalis, M.K. O'Neill, and M.M. West
New Mexico St. Univ. College of Agricultural, Consumer and Environmental Sciences. Agric. Exp. Stn and Coop. Ext. Ser.

Table 12 (cont.). Characteristics and performance of alfalfa varieties across years and tests in New Mexico.		Varietal Characteristics ¹											Las Cruces			Artesia				Tucumcari						Los Lunas		Mora		Farmington								
													2014 ²							2012			2013		2014		2013		2013		2014							
		Variety	Proprietor	RR	FD	WS	Pest resistance								N ³	D	ET	2013	2014	Std			RR [®]			2013	2013	2014										
BW	FW						AN	PRR	SAA	PA	BAA	SN	RKN	15 ⁴	15	15	14	15	14	15	13	14	15	13	14	15	14	15	14	15	15							
Archer III	America's Alfalfa		5	2	HR	HR	HR	HR	n/r	HR	n/r	HR	HR																								*	
Gunner	Croplan Genetics		5	1	HR	HR	HR	HR	HR	R	n/r	R	n/r																									
Mallard	Blue River Hybrids		5	3	HR	HR	HR	HR	R	HR	n/r	R	n/r										*	*													*	
Mountaineer 2.0	Croplan Genetics		5	2	HR	HR	HR	HR	R	HR	n/r	HR	R																								*	
PGI 557	Alforex Seeds		5	2	HR	HR	HR	HR	n/r	R	R	HR	n/r																									
WL 363HQ	W-L Research		5	2	HR	HR	HR	HR	R	R	MR	MR	MR																								*	
Zia	NMSU/Roswell Seed		5	n/r	MR	MR	S	S	MR	S	S	n/r	n/r				*	*	*	*																		
56S82	Pioneer HiBred Int'l		6	5	HR	HR	HR	HR	HR	HR	HR	HR	HR				**	*	*	**	**	**	*	**														
Cimarron VL600	Cimarron USA		6	n/r	R	R	R	HR	HR	HR	R	R	n/r																									
Cisco II	Alforex Seeds		6	2	HR	HR	HR	R	n/r	HR	n/r	R	R	*																								
HybriForce-2600	Sharp Brothers		6	n/r	HR	HR	HR	HR	n/r	n/r	n/r	n/r	n/r																									
Lahontan	USDA/Univ. of Nevada		6	n/r	R	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r																								*	
Transition 6.10RR	Croplan Genetics	Y	6	n/r	R	R	R	HR	R	HR	n/r	MR	n/r																						*			
Wilson	NMSU/Roswell Seed		6	n/r	R	R	n/r	n/r	MR	R	n/r	MR	n/r		*		*	*	*	*	*													*	*	*		
WL 440HQ	W-L Research		6	n/r	HR	HR	R	HR	HR	HR	HR	HR	HR																									
WL 454HQ.RR	W-L Research	Y	6	n/r	R	HR	HR	HR	R	HR	n/r	HR	n/r										*										*	**				
57Q53	Pioneer Hi-Bred Int'l.		7	n/r	MR	HR	HR	R	MR	HR	R	R	MR																									
Artesian Sunrise	Croplan Genetics		7	n/r	MR	HR	R	HR	HR	HR	R	R	n/r				*	*	*	*	*													**	*			
Dona Ana	NMSU/Roswell Seed		7	n/r	MR	MR	LR	R	MR	R	n/r	n/r	n/r				*	*	*	*	*	**	*	*										**	*			
Malone	NMSU/Roswell Seed		7	n/r	R	HR	R	R	R	HR	S	MR	n/r				*	*	*	*	*	**	*	*								*						
NuMex Bill Melton	New Mexico State University		7	n/r	MR	R	R	R	R	MR	MR	n/r	n/r		*				*	*	*	**	*	*											**	*		
SW 7410	S & W Seeds		7	n/r	R	R	HR	MR	HR	R	R	MR	R																									
58N57	Pioneer HiBred Int'l		8	n/r	LR	R	HR	HR	R	HR	HR	MR	HR				*	*	*	*																		
AmeriStand 803T	America's Alfalfa		8	n/r	MR	HR	m	h	r	HR	HR	HR	HR	**	*	*																						
Sandpiper	Blue River Hybrids		8	4	HR	HR	HR	H	R	HR	R	HR	n/r																									
SW 8421S	S & W Seeds		8	n/r	HR	HR	n/r	R	HR	R	R	n/r	R																									
WL 535HQ	W-L Research		8	n/r	n/r	HR	n/r	HR	HR	n/r	n/r	R	R				*	*	*	*																		

¹RR=Roundup Ready if "Y"; WS=Winter Survival (1=No injury, 6=Dead plants), FD=Fall Dormancy (1 & 2 Very dormant; 3 & 4 Dormant; 5 Moderately dormant; 6 & 7 Semi-dormant; 8 & 9 Non-dormant; 10 & 11 Very non-dormant), BW=Bacterial wilt, PRR=Phytophthora root rot, FW=Fusarium wilt, AN=Anthracnose, SAA=Spotted alfalfa aphid, PA=Pea aphid, BAA=Blue alfalfa aphid, SN=Stem nematode, RKN=Rootknot nematode (southern); (S=Susceptible, LR=Low resistance, MR=Moderate resistance, R=Resistant, HR=High resistance, n/r indicates either that the variety was not rated for that characteristic or no rating was available).

²Establishment year.

³N, D, ET, STD, and RR[®] signify normal irrigation, drought irrigation, early termination irrigation, standard performance evaluation, and Roundup Ready[®] entries only, respectively.

⁴Harvest year.

Shaded boxes indicate that the variety was not in the test.

***Significantly higher than all other values in the column.

**Highest yielding variety in the test for that year, except for Los Lunas 2013 test when it is the second highest yielding variety.

*Not significantly different from the highest yielding variety in the test for that year, except for Los Lunas 2013 test in 2014 when it is not significantly different from the second highest yielding variety.

L.M. Lauriault, I.M. Ray, C.A. Pierce, O. Burney, R.P. Flynn, M.A. Marsalis, M.K. O'Neill, and M.M. West
New Mexico St. Univ. College of Agricultural, Consumer and Environmental Sciences. Agric. Exp. Stn and Coop. Ext. Ser.

Table 13. New Mexico State University Agricultural Experiment Station and Cooperative Extension Service publications related to alfalfa management.

Number	Title
A-114	Test your soil
A-122	Soil test interpretations
A-123	Sampling for plant tissue analysis
A-129	Nitrogen fixation by legumes
A-130	Inoculation of legumes
A-131	Certified seed
A-137	Soil analysis: A key to soil nutrient management
A-145	Certified noxious weed free program
A-229	Phymatotrichum root rot
A-325	Managing weeds in alfalfa
A-326	Downy mildew on alfalfa
A-333	User manual of the alfalfa yield predictor
A-334	Beet armyworm in New Mexico Hay
A-335	Variegated cutworm in New Mexico Hay
A-336	Managing Roundup Ready alfalfa and conventional or organic alfalfa hay in nearby fields in New Mexico
A-337	Recommendations for Roundup Ready alfalfa weed management and stand removal in New Mexico
A-338	Alfalfa weevil control options in New Mexico
A-339	Alfalfa integrated pest management: Aphids
H-158	How to collect and send plant specimens for disease diagnosis
CR-536	Blister beetles in alfalfa
CR-633	Using a computer application to predict irrigated alfalfa yield
CR-641	Hay quality, sampling and testing
CR-644	Assessing alfalfa stands after winter injury, freeze damage, or any time renovation is considered in New Mexico
CR-646	Managing alfalfa during drought
CR-654	Selecting alfalfa varieties for New Mexico
CR-659	Whitefringed beetle in New Mexico alfalfa
CR-668	Reducing harvest and post-harvest losses of alfalfa and other hay
RR-766	Furrow-irrigated alfalfa dry matter yield is not affected by different seeding rates in the Southern High Plains, USA
RR-772	Observations on how cowpea aphid affects alfalfa

These publications, and alfalfa variety test reports from previous years, are available from your county office of the NMSU Cooperative Extension Service or online at <http://forages.nmsu.edu/resources.html> and aces.nmsu.edu/pubs/