



The 2002 New Mexico Alfalfa Variety Test Report



Agricultural Experiment Station
College of Agriculture and Home Economics

The 2002 New Mexico Alfalfa Variety Test Report

L.M. Lauriault, I.M. Ray, D.A. McWilliams, L.M. English, R.P. Flynn, S.J. Guldán, M.W. Murray, and M.K. O'Neill¹

Introduction

In 2002, 260,000 acres of alfalfa (*Medicago sativa*) remained in production in New Mexico yielding an estimated 1.3 million tons of hay. Because of prolonged drought, hay stocks in New Mexico and throughout the Southwest are dwindling, resulting in higher prices received. At an average of \$138/ton (up from \$126 in 2001), gross returns will total approximately \$183 billion, ensuring that alfalfa hay remains New Mexico's No. 1 cash crop (New Mexico Agricultural Statistics Service Weekly AgUpdate). Alfalfa also is the legume of choice in irrigated perennial pastures.

Choosing a good alfalfa variety is a key step in establishing a highly productive stand of alfalfa whether for hay or pasture. Differences between the highest- and lowest-yielding varieties in tests included in this report ranged from 0.49 to 6.68 tons per acre in 2001. If sold as hay, this translates to a difference in returns of \$62 to \$842 per acre due to variety. Stand longevity, as affected by winter hardiness and pest resistance, also is partially determined by variety.

New trends in alfalfa varieties. In recent years a number of alfalfa varieties have been developed for tolerance to frequent defoliation and hoof damage under grazing. True grazing-tolerant varieties have a broad crown set below the soil surface. This trait gives protection from hoof damage and permits retention of greater root carbohydrate reserves for continued growth. The deep-set crown also helps prevent damage by other factors, such as equipment traffic. Grazing-tolerant alfalfa varieties also have the ability to produce and retain leaves below the grazing horizon, which allows photosynthesis to continue even under frequent defoliation of upper leaves. The development of grazing tolerance was accomplished using continuous grazing during the growing season. However, in New Mexico, many alfalfa fields are used for overwintering pastures and the effects of this practice have not yet been fully measured.

Another beneficial development for alfalfa producers is subsurface irrigation, or drip irrigation. This irrigation technique lessens evaporation and runoff losses and might allow for an increase in irrigated acreage. Previously used for high-value annual crops, such as vegetables, problems with roots and salts clogging emitters prevented use with perennial crops. Improvements in tubing, flushing systems and management practices might allow drip irrigation to be used

in alfalfa.

This report, which is a collaborative effort of New Mexico State University scientists at agricultural science centers throughout the state, provides yield data for alfalfa varieties included in yield trials in New Mexico and guidelines for variety selection.

Considerations in Selecting an Alfalfa Variety

Local adaptation and persistence. High yields in variety tests over a number of years and locations within a region are the best indication that a variety is locally adapted and persistent. Select varieties based on unbiased information collected from trials conducted by NMSU scientists. Look first at data collected from the agricultural science center(s) closest to you. Data from the centers is grouped by similarity of latitude and elevation, which affect winter hardiness and, thus, yield and persistence.

Persistence is the ability of individual plants to survive field conditions over time and is strongly influenced by winter hardiness, harvest frequency, and pest resistance. Higher persistence will permit a longer stand life in which to recover establishment costs. Alfalfa stands should be replaced when plant density drops to <5 plants (or 40 stems) per square foot. Producers should rotate to another irrigated crop for at least one year before reseeding alfalfa into the same field to avoid seedling death due to autotoxicity. If stand life expectancy is only 3 to 4 years, as is the case in a crop rotation system, higher yields in those early years are more important than persistence.

Winter hardiness. Alfalfa's winter hardiness is determined by its ability to survive cold temperatures. In the past, winter hardiness was given by the fall dormancy rating (FD), which indicates the variety's tendency to stop growing in the fall. Fall dormancy categories range from 1 (very dormant) to 9 (nondormant). The more dormant (FD 1 to 3) varieties will be slower to "green up" in the spring and quicker to stop growing in the fall, regardless of local climate. This will have an impact on yield in areas with warmer climates. Additionally, nondormant varieties (FD 7 to 9) tend to yield more because of earlier spring "green up" and later fall production but might not survive severe winters. These varieties might be suitable for use in a short-term rotation system, where alfalfa is used for 4 years or less. Otherwise, produc-

¹Forage agronomist, NMSU Agricultural Science Center at Tucumcari; alfalfa breeder, NMSU, Las Cruces; Extension agronomist, NMSU, Las Cruces; superintendent, NMSU Agricultural Science Center at Los Lunas; superintendent, NMSU Agricultural Science Center at Artesia; superintendent, NMSU Sustainable Agriculture Science Center at Alcalde; assistant professor, NMSU Agricultural Science Center at Artesia; and superintendent, NMSU Agricultural Science Center at Farmington, respectively.

ers should select varieties with sufficient dormancy to survive winter conditions at their location, while optimizing forage production during the growing season.

The North American Alfalfa Improvement Council has developed another classification for winter hardiness, Winter Survival (WS), because some varieties are more able to withstand low temperatures than their fall dormancy category indicated. This system rates varieties from 1 to 6, where a rating of 1 indicates little or no winter injury and 6 indicates plant death. Ratings are made after the first winter for spring seedlings. At least two location years are required for the standard test. Because these tests are conducted in areas that have severe winters, only varieties in fall dormancy categories 6 or less currently are being rated for winter survival.

Variety test reports, such as this one, also are valuable tools for determining the true winter survivability of any alfalfa variety, particularly in the northern half of New Mexico and the north-central mountains where several nondormant (FD 7-9) varieties have performed well. When using the winter survival data, keep in mind that these tests usually are conducted in the northern states, and varieties that will not survive their more severe winters will likely survive in New Mexico.

Disease resistance. New Mexico alfalfa producers should select adapted varieties with the highest available resistance, preferably an “R” (resistant) or greater rating to bacterial wilt (Bw), Fusarium wilt (Fw), Phytophthora root rot (PRR), and anthracnose (An) (Alfalfa Analyst, Certified Alfalfa Seed Council).

Bacterial and Fusarium wilt are water-conducting tissue infections of alfalfa’s roots that do not cause any noticeable root rot. These diseases prevent water flow to leaves, resulting in wilted shoots and, eventually, death of infected plants. Roots infected with bacterial wilt often will have a yellowish brown discoloration of the taproot’s inner woody cylinder. Fusarium infection, on the other hand, is recognized by brown to red streaks in the taproot’s inner woody cylinder.

Phytophthora root rot is a fungal disease associated with excessive soil moisture. This disease causes yellowish to brown areas on roots and crowns that eventually become black and rotten. Top growth of infected plants appears stunted and yellow.

Anthracnose, also caused by a fungus, attacks alfalfa stems, preventing water flow to the rest of the shoot and causing sudden wilting. These wilted shoots have a characteristic “shepherd’s crook” appearance. Anthracnose also can cause a bluish black crown rot.

Many other alfalfa diseases also occur in New Mexico, resistance to which has not yet been developed. The best protection against them is proper management. But as producers know, even that, at times, is not sufficient.

Insect resistance. There are many insects that feed on alfalfa. Varietal resistance is available for spotted alfalfa aphid (SAA), pea aphid (PA) and blue alfalfa aphid (BAA), which have been the most detrimental insects to alfalfa production in New Mexico. As with disease resistance, select

varieties that have at least an “R” rating for each of the insects. Varieties with resistance to insects that are not adapted to your area might not be preferred, due to a historical decrease in yield associated with the resistance.

Another insect, cowpea aphid, has begun infesting alfalfa fields in New Mexico in recent years. Adult cowpea aphids are smaller than other common aphids. They are black with white or yellow markings on legs and antennae. Nymphs are gray to purple and can be confused with blue alfalfa aphid. Critical infestations can occur in early spring when alfalfa breaks dormancy. Plants are severely stunted, reducing first cutting yields and causing possible stand loss if not controlled.

Currently no resistant varieties have been released, but they are being developed. Predatory insects can help control cowpea aphid populations. Unfortunately, the predators might not be active in early spring when the alfalfa greens up. Chemical control might be the only option in early spring. Economic thresholds and monitoring procedures have not been established. Scout several areas in each field for the presence of aphids (purple or black) and stunted plants. When infestations are found, immediate treatment is imperative to prevent further damage and stand loss. Contact your county Cooperative Extension Service office or a licensed pesticide applicator about what products can be used to control cowpea aphids. Even if infestation is found in only one field, it might be advisable to spray all fields to prevent migration or to control infestations previously undetected.

Varietal resistance to other insects, such as alfalfa weevil and potato leafhopper, also might be available in the near future. However, historic resistance has not protected the plant. Rather it masks the symptoms. Currently, the best protection against these insect pests is proper harvest management and pesticide use.

Seed quality. Selecting an alfalfa variety based on seed cost is like playing Russian roulette. Seed labeled “common,” “variety not stated,” or “variety unknown” are of unknown genetic background and may or may not be locally adapted or have the necessary disease or insect resistance. To be assured of achieving a long-lasting, highly productive stand, buy either certified or Plant Variety Protected (PVP) seed, which guarantees the genetics and performance. Look for a blue tag, which must be attached to all bags of certified seed, or Plant Variety Protection labeling, which is the proprietor’s guarantee. Be sure to read the seed tag, which provides important information about the seed, including purity, amounts of other crop and weed seed (including any noxious weed seed), germination, and the test date (within the previous 9 months). Order seed well in advance of planting time to assure that it will be available when needed.

Forage quality. High quality alfalfa hay possesses the following characteristics: >19 percent crude protein, <31 percent acid detergent fiber, <40 percent neutral detergent fiber, leafiness, and free of foreign material. Varietal differences in quality are relatively small compared to other factors. Cultural and management practices, such as soil fertil-

ity, irrigation, weed and insect control, maturity at cutting, baling, and storage conditions, are major factors that affect alfalfa quality.

Table 1 gives quality data for alfalfa varieties in a test sown in 2001 at Farmington. No differences existed between varieties across fall dormancy categories, but there are differences between fall dormancy categories. Standard Deviations (SD) of variety means are included for each quality variable. The SD measures the variability between different samples of the same variety. For example, each variety in the 2001 Farmington test was sampled (replicated) four times. The crude protein (CP) of any of the four samples for any variety can vary from the average for that variety by as much as the associated SD. The lack of difference between varieties is demonstrated, in that, the largest SD given for any variable is nearly equal to the difference between the means of the highest and lowest varieties for each variable.

Differences in quality between fall dormancy categories are likely due to stage of maturity at harvest. Recovery after harvest is faster as fall dormancy category increases from 1 to 9 (from dormant to nondormant). Therefore, if varieties in different fall dormancy categories always are harvested on the same day, as is the case for most variety tests, those in lower fall dormancy categories usually will be at an earlier stage of maturity than those in higher categories. Harvesting earlier might give lower yields, but it also will likely give higher quality.

The optimum balance between forage yield, quality, and plant persistence occurs at 1-10 percent bloom. Harvesting at prebloom increases quality but sacrifices yield. Continued harvesting at prebloom reduces stand life, because the plant is not able to replenish root reserves for subsequent growth and overwintering. Since some hay buyers specify prebloom, producers should weigh price against decreased yields and shorter stand life. Prebloom harvests in the middle cuttings are likely to be less detrimental to stand life than the first and last cuttings. Cutting at less than 10 percent bloom increases yield, but quality declines rapidly as fiber increases and mineral content decreases (Alfalfa for Dairy Animals, Certified Alfalfa Seed Council). Insect feeding, maturity and harvesting affect leaf retention and, therefore, forage quality, because the digestibility and nutrient content in leaves is greater than in stems.

Description of Tests

Replicated alfalfa variety tests included in this report were conducted under research controls at NMSU's agricultural science centers at Las Cruces (sown in 1999), Artesia (1999, 2000 and 2001), Tucumcari (1999 and 2001), Los Lunas (1999 and 2001), Alcalde (2001) and Farmington (1999 and 2001). In addition to standard yield trials sown in 2001, new tests this year include a grazing tolerance trial sown in 2001, at Alcalde that will begin comparing the effects of winter grazing, after the winter of 2002-2003; and a test sown in 2000 at Artesia under drip irrigation.

Weather data for 2001 and the long-term averages

from all locations are presented in table 2. Yield data (on a dry matter basis) are presented in tables 3-13. Varieties are listed in order from highest to lowest average annual production. Yields are given by cutting for 2001 and by year for each production year. Statistical analyses were performed on all alfalfa yield data (including experimentals) to determine if the apparent differences are truly due to variety or just to chance (SAS). The variety with the highest numerical yield in each column is marked with two asterisks (**), and those varieties not significantly different from that variety are marked with one asterisk (*). To determine if two varieties are truly different, compare the difference between the two varieties to the Least Significant Difference (LSD) at the bottom of the column. If the difference is equal or greater than the LSD, the varieties are truly different in yield when grown under the conditions at a given location. The Coefficient of Variation (CV), which is a measure of the variability of the data, is included for each column of means. Low variability (<20 percent) is desirable, and increased variability within a study results in higher CVs and larger LSDs. There might be a difference between previously published data and the data given in this publication for the same tests because of differences in the programs used for statistical analysis.

Table 14 summarizes information about proprietors, winter survival (measured in the northern United States), fall dormancy, pest resistance and yield performance across years and locations for all varieties currently included in NMSU's alfalfa variety testing program. Varieties are listed alphabetically by fall dormancy category. Be cautious in using the winter survival data, because it might underestimate a variety's winter survivability in New Mexico. Long-term performance in the northern half of the state should be a good indication of winter hardiness. In table 14, shaded areas indicate that the variety was not in that particular test (labeled at the top of the column), while clear blocks mean that the variety was in the test. As before, a double asterisk (**) indicates that the variety had the highest yield in the test for that year, and a single asterisk (*) means that the variety was not significantly different from the highest-yielding variety based on the 5 percent LSD. It is best to choose a variety that has performed well over several years and locations as indicated by the asterisks. Tests are grouped by location, which are grouped by similar elevations and latitudes within New Mexico.

Once varietal choices are made using table 14, look at cutting data (tables 3 to 14) to make sure the variety will be productive during the desired season. Varieties selected for grazing should produce over a longer season. Those used for hay should produce well in times that avoid potential problems. For instance, horse hay should be harvested early in the season to avoid blister beetle infestations that might occur in the later cuttings. Higher-value dairy hay might be produced later in the season to avoid spring weed problems.

Regarding "common" varieties, notice that the same variety might yield well in one area but not another. And those that do yield well might not do so consistently across years. Generally, those that produce well will do so until a pest

problem occurs. Then the stand can be lost, requiring a waiting period before reseeding.

Summary

Consistent production of high alfalfa yields is the result of selecting good varieties, along with implementing good management techniques. Soil fertility should be maintained at recommended levels based on soil tests, and weeds and insects should be controlled using appropriate cultural and/or chemical methods. For dormant (FD 1 to 3) and semidormant (FD 4 to 6) varieties, a 6-week rest period before a dormancy-inducing freeze (27°F) is recommended to allow plants to replenish root reserves for winter survival and initiate spring growth, after which harvesting might be done either mechanically or by grazing. Removing fall growth is beneficial to reducing weevil populations the following year as eggs are laid in and overwinter in stems. Harvesting established stands at early bloom would result in 3 to 5 cuttings per year before initiation of the rest period in most of New Mexico. For further information about alfalfa management, refer to the other NMSU Cooperative Extension Service publications listed in table 15.

Table 1. Forage quality¹ of alfalfa varieties sown August 2001, at the NMSU Agricultural Science Center at Farmington, 2002².

Variety	Fall dormancy	CP, %		ADF, %		NDF, %		RFV		Fall dormancy means			
		Mean	SD	Mean	SD	Mean	SD	Mean	SD	CP, %	ADF, %	NDF, %	RFV
Champ	3	22.03*	±0.52	29.30*	±0.99	34.45*	±1.02	179.08*	±6.43	22.03**	29.30*	34.45	179.08*
DKA42-15	4	21.57*	±0.72	27.83*	±0.98	33.32*	±1.52	188.73**	±10.27				
WL327	4	22.17*	±1.13	28.36*	±0.80	33.61*	±2.32	187.11*	±15.76				
Megaton 3.5	4	22.06**	±1.68	28.05*	±0.94	33.85*	±1.25	184.82*	±8.88				
Geneva	4	21.50*	±1.09	28.41*	±0.92	33.77*	±1.17	184.68*	±7.91				
Legend	4	21.38*	±0.97	28.31*	±1.46	34.09*	±1.30	183.27*	±9.70				
Delta526	4	21.58*	±1.12	29.10*	±1.85	33.86*	±1.36	182.80*	±10.00				
Focus HSN	4	21.31*	±0.61	29.20*	±1.16	34.23*	±1.88	181.04*	±12.30				
Magnum V	4	21.36*	±0.76	28.52*	±0.56	34.44*	±0.95	180.28*	±4.93				
Forecast 1001	4	21.16*	±0.85	29.48*	±1.31	34.51*	±0.95	177.95*	±7.51				
HybriGreen 41	4	21.12*	±1.53	29.21*	±0.62	34.88*	±0.85	176.63*	±4.09				
WL342	4	20.98*	±1.00	28.78*	±0.63	35.24*	±1.49	176.20*	±5.51				
54V54	4	20.89*	±0.46	29.41*	±0.91	35.66*	±0.81	172.27*	±5.25				
HybriForce 400	4	20.54*	±0.43	30.12*	±1.53	35.74*	±1.31	170.61*	±9.07	21.36*	28.83*	34.40	180.49**
5-Star	5	21.36*	±0.40	28.81*	±0.59	33.92*	±1.15	182.88*	±7.35				
Archer II	5	20.92*	±0.33	29.72*	±1.18	35.50*	±0.68	172.46*	±5.29	21.14*	29.26*	34.71	177.67*
Dona Ana	8	19.95*	±0.61	30.55**	±0.91	37.33*	±0.72	162.48*	±4.55	19.95	30.55**	37.33*	162.48
NM-9D11A-PAR	?	20.83*	±1.80	29.54*	±1.22	37.65**	±1.87	163.22*	±8.07	20.83	29.54*	37.65**	163.22
Mean		21.26		29.04		34.78		178.14					
Variety LSD, 0.05		Ns		Ns		Ns		Ns		1.28	Ns	1.79	11.85
CV, %		4.21		3.55		3.60		4.65					

¹Appreciation is expressed to the Navajo Agricultural Products Industry Research and Testing Laboratory for conducting the analysis.

²Data are the means of two harvests taken 18 July and 27 August 2002.

CP, ADF, NDF, RFV, SD, LSD, Ns, and CV signify crude protein, acid detergent fiber, neutral detergent fiber, relative feed value, standard deviation, least significant difference, not significantly different based on a protected statistical analysis, and coefficient of variation, respectively. The SD gives an indication of the variability was found within a variety. Generally, individual observations (4 replicates for each variety for the data presented in this table) can be expected vary from the mean by as much as the SD. If the difference between means within a column is equal to or greater than the LSD given at the bottom of that column, we are 95% certain that they are truly different. The CV gives an indication of the amount of variation accounted for in the statistical analysis of a variable. Sources of variation included in the statistical model for these data include replicate, fall dormancy, and variety.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

Table 2. Temperature and precipitation data for 2002 and the long-term averages for the New Mexico alfalfa variety test locations.

Location	Las Cruces ¹				Artesia				Tucumcari				Los Lunas				Alcalde				Farmington			
Elevation	3832 ft.				3376 ft.				4091 ft.				4842 ft.				5725 ft.				5577 ft.			
Latitude	32° 12' N				32° 45' N				35° 12' N				34° 16' N				36° 05' N				36° 41' N			
Month	Temp. (°F)		Precip. (In)		Temp. (°F)		Precip. (In)		Temp. (°F)		Precip. (In)		Temp. (°F)		Precip. (In)		Temp. (°F)		Precip. (In)					
	02	Ave.	02	Ave.	02	Ave.	02	Ave.	02	Ave.	02	Ave.	02	Ave.	02	Ave.	02	Ave.	02	Ave.				
Nov-01	51	50	0.19	0.53	51	48	0.95	0.50	52	47	1.57	0.63	47	49	1.14	0.45	43	40	0.35	0.72	31	41	0.48	0.71
Dec-01	40	42	0.20	0.68	41	40	0.03	0.47	42	39	0.24	0.58	34	35	0.12	0.51	30	31	0.03	0.43	31	32	0.55	0.42
Jan-02	42	42	0.00	0.56	42	39	0.24	0.52	41	38	0.54	0.34	36	44	0.49	0.34	31	30	0.83	0.50	32	30	0.04	0.50
Feb-02	45	46	1.27	0.37	42	44	0.55	0.41	41	42	0.16	0.47	37	44	0.05	0.46	31	36	0.09	0.42	33	36	0.04	0.42
Mar-01	51	52	0.00	0.22	49	51	1.33	0.37	47	48	0.32	0.68	46	54	0.00	0.50	41	43	0.00	0.67	41	43	0.17	0.73
Apr-01	65	59	0.00	0.21	64	60	0.02	0.50	60	57	0.53	1.13	39	52	0.13	0.46	54	51	0.30	0.61	57	50	0.37	0.59
May-02	68	68	0.31	0.29	70	68	0.00	1.20	70	65	0.65	1.98	66	64	0.19	0.46	60	60	0.00	0.55	62	60	0.00	0.55
Jun-01	76	77	0.00	0.72	81	75	0.00	1.54	81	75	0.67	1.83	76	73	0.08	0.56	70	70	1.37	0.29	75	70	0.00	0.27
Jul-02	81	80	0.32	1.36	80	79	0.00	1.50	80	79	4.38	2.61	77	78	1.35	1.24	73	75	0.67	0.89	78	76	0.42	0.88
Aug-02	77	78	1.63	2.29	81	77	0.00	2.12	81	77	0.72	2.69	76	76	0.62	1.80	70	74	0.29	1.11	74	74	0.32	1.12
Sep-02	77	72	0.46	1.38	72	70	0.00	2.11	71	70	4.00	1.42	68	66	3.05	1.30	63	66	3.72	1.08	61	66	3.26	1.03
Oct-02	67	61	0.40	0.91	60	59	2.32	1.19	57	59	1.20	1.29	57	69	0.76	0.97	51	53	1.53	0.92	52	54	1.75	0.93
Annual	62	61	4.78	9.40	61	59	5.44	12.43	60	58	14.98	15.65	55	59	7.98	9.05	51	52	9.18	8.15	54	52	7.40	8.16

¹Long-term averages for the Las Cruces test site are from the State University weather station, located approximately 5.5 miles to the north.

Table 3. Dry matter yields (tons/acre) of alfalfa varieties sown September 8, 1999, at NMSU's Leyendecker Plant Science Research Center at Las Cruces and flood-irrigated every 14 days.

Variety	2000 Total ¹	2001 Total	2002 Harvests						2002 Total	3-yr Av- erage
			30-Apr	10-Jun	15-Jul	13-Aug	27-Sep	7-Nov		
CW 68115	13.83*	10.72**	2.05*	2.10*	2.29**	1.33*	1.47**	1.41**	10.65**	11.73**
CW 5875	13.87*	10.54*	2.00*	1.98*	2.24*	1.36**	1.47**	1.38*	10.42*	11.61*
CW 78122	13.50*	10.25*	2.04*	2.01*	2.26*	1.29*	1.45*	1.38*	10.43*	11.39*
CW 5666	13.45*	10.50*	2.09*	1.89*	2.04*	1.21*	1.43*	1.27	9.94*	11.30*
NM9D11A-PAR	13.92**	9.80*	2.11**	2.13**	2.12*	1.21*	1.37*	1.16	10.10*	11.27*
GT 13-R Plus	13.09*	10.16*	1.94*	1.89*	2.04*	1.26*	1.46*	1.30*	9.89*	11.05*
Rio Grande	13.63*	9.91*	1.98*	1.84*	1.84	1.04	1.28	1.20	9.16	10.90*
NM stress 94	13.34*	9.84*	2.05*	1.99*	1.82	0.95	1.32*	1.19	9.32*	10.84*
NM Common	13.50*	9.35	1.83*	1.75*	1.92*	1.20*	1.44*	1.17	9.31*	10.72*
Mesa	13.11*	9.69*	1.90*	1.77*	1.88*	1.14*	1.42*	1.21	9.32*	10.71*
Arriba	13.14*	9.81*	1.91*	1.73*	1.79	1.13	1.39*	1.18	9.13	10.69*
AmeriLeaf 721	13.36*	9.71*	1.98*	1.61	1.69	1.00	1.29	1.15	8.72	10.60*
57Q77	13.02*	9.48	1.94*	1.72*	1.76	1.01	1.35*	1.30*	9.07	10.52*
PGI 8000	12.89*	9.65*	1.81	1.72*	1.87*	1.03	1.31*	1.21	8.94	10.50*
Signal 8000	12.73	9.64*	1.85*	1.81*	1.77	1.06	1.27	1.24	9.00	10.46*
Mesa	13.22*	9.13	1.79	1.74*	1.87*	1.00	1.36*	1.24	9.00	10.45
Wilson	12.35	9.37	2.11**	1.92*	1.95*	1.11	1.31*	1.07	9.46*	10.39
DS 482	13.47*	8.85	1.69	1.52	1.83	1.04	1.32*	1.18	8.57	10.30
ZG9891	12.49	9.47	1.96*	1.62	1.66	0.93	1.27	1.13	8.57	10.18
WL 612	12.90*	9.16	1.59	1.56	1.67	1.05	1.30	1.24	8.42	10.16
ZX9889B	13.42*	8.88	1.74	1.53	1.68	0.92	1.12	1.18	8.17	10.16
AmeriGraze 701	12.73	9.34	1.90*	1.50	1.58	0.94	1.21	0.96	8.09	10.05
CutMor	12.86*	8.87	1.63	1.59	1.85	0.98	1.24	1.14	8.42	10.05
DK180ML	12.42	9.01	1.69	1.73*	1.74	0.96	1.29	1.18	8.60	10.01
Dona Ana	11.83	8.94	1.85*	1.71*	1.77	1.04	1.23	1.08	8.68	9.82
WL 525HQ	12.21	8.65	1.66	1.55	1.59	1.08	1.28	1.25	8.40	9.76
WL 442	13.20*	8.56	1.68	1.39	1.42	0.82	1.14	1.00	7.44	9.73
ZL9876	12.50	8.56	1.78	1.47	1.42	0.85	1.19	1.14	7.86	9.64
ZX9894	11.87	8.67	1.74	1.50	1.64	0.95	1.22	1.20	8.25	9.60
58N57	10.56	7.94	2.02*	1.39	1.38	0.84	1.14	0.81	7.57	8.69
Mean	12.95	9.42	1.88	1.72	1.81	1.06	1.31	1.19	8.96	10.44
LSD (0.05)	1.17	1.19	0.30	0.43	0.43	0.23	0.17	0.13	1.43	1.28
CV%	6.41	9.01	11.27	17.59	16.91	15.65	8.97	7.82	11.33	13.50

¹Yield data from previous years may be different than that presented in other publications due to a difference in analysis methods.

2000 Harvest dates: 3-May, 1-Jun, 7-Jul, 4-Aug, 16-Sep, and 2-Nov.

2001 Harvest dates: 1-May, 11-Jun, 13-Jul, 14-Aug, 27-Sep, and 7-Nov.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

LSD (0.05) stands for the Least Significant Difference at the 5% level. If the difference between two numbers within a column is equal to or greater than the LSD, it is 95% certain that they are truly different.

Ns means that there were no significant differences between varieties within that column at the 5% level.

Table 4. Dry matter yields (tons/acre) of alfalfa varieties sown September 2, 1999, at NMSU's Agricultural Science Center at Artesia and flood-irrigated approximately twice per cutting.

Variety	2000 Total	2001 Total	2002 Harvests						2002 Total	3-yr Average
			7-May	2-Jun	3-Jul	30-Jul	3-Sep	8-Nov		
Artesian Sunrise	8.69**	8.06*	1.80*	1.75*	1.93*	1.25*	0.88*	0.39*	7.55*	8.10**
AmeriStand 802	7.87*	8.35**	1.93**	1.83*	1.85*	1.35*	0.93*	0.43*	7.93**	8.05*
AmeriStand 801S	7.96*	8.21*	1.83*	1.75*	1.88*	1.43**	1.05**	0.44*	7.90*	8.02*
57Q77	8.40*	7.90*	1.90*	1.88**	1.70*	1.23*	1.00*	0.44*	7.70*	8.00*
DS 771	8.13*	7.85*	1.88*	1.80*	1.90*	1.15*	0.90*	0.40*	7.60*	7.86*
DS 981	7.76*	8.18*	1.83*	1.65*	1.80*	1.33*	0.98*	0.44*	7.60*	7.85*
AmeriGraze 701	7.82*	7.84*	1.88*	1.73*	1.93*	1.23*	0.85*	0.42*	7.60*	7.75*
58N57	8.58*	7.47*	1.58*	1.65*	1.73*	1.20*	0.90*	0.39*	7.05*	7.70*
Magna 901	7.65*	7.81*	1.78*	1.75*	1.88*	1.33*	0.90*	0.42*	7.63*	7.69*
ZX 9393	7.30*	8.06*	1.65*	1.68*	2.00**	1.33*	1.03*	0.41*	7.63*	7.66*
WL 442	7.70*	7.90*	1.48*	1.68*	1.88*	1.35*	0.95*	0.42*	7.30*	7.63*
Rio Grande	7.48*	7.90*	1.85*	1.75*	1.80*	1.20*	0.88*	0.41*	7.48*	7.62*
NM9D11A-PAR	7.21*	7.79*	1.80*	1.75*	1.93*	1.33*	1.03*	0.45**	7.83*	7.61*
WL 525HQ	7.53*	7.51*	1.68*	1.75*	1.95*	1.35*	1.00*	0.43*	7.73*	7.59*
CW 5875	7.40*	7.95*	1.78*	1.68*	1.73*	1.30*	0.90*	0.42*	7.40*	7.58*
CW 68115	7.47*	8.00*	1.58*	1.73*	1.78*	1.28*	0.90*	0.42*	7.20*	7.56*
CW 78122	7.01*	8.15*	1.85*	1.73*	1.80*	1.18*	0.88*	0.37*	7.45*	7.54*
CutMor	7.59*	7.46*	1.65*	1.75*	1.98*	1.28*	0.88*	0.44*	7.50*	7.52*
DK180 ML	7.12*	7.89*	1.78*	1.63*	1.83*	1.30*	0.93*	0.44*	7.50*	7.50*
Tahoe	7.25*	7.68*	1.88*	1.75*	1.85*	1.20*	0.88*	0.41*	7.48*	7.47*
WL 414	7.80*	7.32*	1.65*	1.60*	1.83*	1.20*	0.93*	0.41*	7.18*	7.43*
CW 5666	7.48*	7.50*	1.78*	1.68*	1.80*	1.18*	0.90*	0.38*	7.30*	7.43*
Arriba	8.10*	7.20*	1.73*	1.63*	1.60*	1.05*	0.85*	0.37*	6.88*	7.39*
DS 681FQ	6.85*	7.67*	1.60*	1.63*	1.88*	1.28*	0.98*	0.38*	7.38*	7.30*
Mean	7.67	7.82	1.76	1.72	1.84	1.26	0.93	0.41	7.49	7.66
LSD (0.05)	Ns	Ns	Ns	Ns	Ns	Ns	Ns	Ns	Ns	Ns
CV%	15.08	7.64	11.20	10.95	12.56	12.94	11.14	14.36	8.57	12.24

¹Yield data from previous years may be different than that presented in other publications due to a difference in analysis methods.

2000 Harvest dates: 3-May, 14-Jun, 20-Jul, and 21-Aug.

2001 Harvest dates: 10-May, 11-Jun, 9-Jul, 7-Aug, 10-Sep, and 29-Oct.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5%LSD.

LSD (0.05) stands for the Least Significant Difference at the 5% level. If the difference between two numbers within a column is equal to or greater than the LSD, it is 95% certain that they are truly different.

Ns means that there were no significant differences between varieties within that column at the 5% level.

Table 5. Dry matter yields (tons/acre) of alfalfa varieties sown September 13, 1999, at NMSU's Agricultural Science Center at Artesia and drip-irrigated.

Variety	Total annual yields			
	2000	2001	2002	3-yr Average
Rio Penasco	8.30*	8.73*	7.74**	8.25**
Rio Grande	8.47*	8.76**	7.18*	8.14*
WL525HQ	8.13*	8.60*	7.26*	8.00*
Signal 8000	8.25*	8.32*	6.97*	7.85*
Dona Ana	8.49**	8.18*	6.57*	7.74*
WL414	8.22*	7.81*	7.06*	7.70*
NM Common	7.90*	8.36*	6.62*	7.62*
Arriba	8.11*	7.47*	7.29*	7.62*
Signal 7000	8.01*	7.70*	6.84*	7.52*
Tahoe	8.27*	7.79*	6.21*	7.42*
Magna901	7.93*	8.05*	6.18*	7.38*
Salado	8.38*	7.89*	5.80*	7.36*
AmeriGraze 702	7.51*	7.81*	6.70*	7.34*
Rio	7.56*	7.17*	6.47*	7.07*
Mean	8.06	8.16	6.78	7.67
CV, %	7.91	11.81	20.30	10.82
DMRT (0.05)	Ns	Ns	Ns	Ns

2000 Harvest dates: 5-May, 13-Jun, 20-Jul, 18-Aug, and 13-Sep.

2001 Harvest dates: 19-Apr, 24-May, 20-Jun, 16-Jul, 17-Aug, and 28-Sep.

2002 Harvest dates: 14-May, 17-Jun, 12-Jul, 15-Aug, and 23-Sep.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% DMRT.

DMRT (0.05) stands for Duncan's Multiple Range Test for differences among all entries at the 5% level. If the difference between two numbers within a column is equal to or greater than the DMRT, it is 95% certain that they are truly different.

Ns means that there were no significant differences between varieties within that column at the 5% level

Table 6. Dry matter yields (tons/acre) of flood-irrigated alfalfa varieties sown September 28, 2001, at NMSU's Agricultural Science Center at Artesia and flood-irrigated approximately twice per cutting.

Variety	2002 Harvests						2002 Total
	15-May	Jun-02	11-Jul	23-Aug	30-Sep	8-Nov	
DS 8181	1.63*	1.50*	2.94*	2.28*	1.55*	0.96	10.86**
Aspire	1.73**	1.55*	2.88*	2.31*	1.52*	0.82	10.81*
AmeriStand 802	1.55	1.44	2.91*	2.28*	1.60*	0.96	10.74*
58N57	1.52	1.49*	2.92*	2.25	1.60*	0.94	10.70*
Tru Test	1.55	1.39	2.88*	2.39**	1.67**	0.81	10.68*
DS 7117	1.65*	1.57**	3.00**	2.25	1.46*	0.75	10.67*
AL 299	1.59*	1.49*	2.89*	2.26	1.57*	0.86	10.65*
Magna 901	1.38	1.37	2.78	2.31*	1.65*	1.09**	10.57*
Duran 2-2	1.58	1.42	2.78	2.39**	1.55*	0.77	10.48*
WL 442	1.52	1.43	2.81	2.29*	1.54*	0.85	10.44*
Magna 601	1.53	1.44	2.77	2.27*	1.52*	0.76	10.28
Pershing	1.36	1.35	2.72	2.21	1.55*	1.01*	10.21
NM9D11A-PAR	1.54	1.47*	2.75	2.33*	1.43*	0.68	10.19
WL 625HQ	1.40	1.40	2.61	2.18	1.54*	1.03*	10.15
WL 525HQ	1.44	1.41	2.61	2.27*	1.44*	0.95	10.11
AmeriStand 801S	1.48	1.39	2.58	2.22	1.54*	0.89	10.08
ZS 9992	1.39	1.33	2.59	2.28*	1.45*	0.92	9.96
ZS 0000	1.32	1.38	2.71	2.15	1.54*	0.83	9.92
CutMor	1.29	1.37	2.64	2.20	1.47*	0.91	9.87
Dona Ana	1.30	1.31	2.65	2.30*	1.43*	0.85	9.84
57Q77	1.50	1.36	2.62	2.18	1.48*	0.69	9.83
NM Common	1.31	1.32	2.58	2.14	1.45*	0.72	9.51
Mean	1.48	1.42	2.76	2.26	1.53	0.87	10.30
LSD (0.05)	0.15	0.12	0.19	0.13	Ns	0.12	0.58
CV%	7.14	6.14	4.93	4.07	8.41	9.97	3.98

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

LSD (0.05) stands for the Least Significant Difference at the 5% level. If the difference between two numbers within a column is equal to or greater than the LSD, it is 95% certain that they are truly different.

Ns means that there were no significant differences between varieties within that column at the 5% level.

Table 7. Dry matter yields (tons/acre) of alfalfa varieties sown September 14, 1999, at NMSU's Agricultural Science Center at Tucumcari and previously furrow-irrigated once per cutting but only on May 15 in 2002 due to water shortages¹.

Variety	2000 Total ²	2001 Total	Yield (dry tons/acre)						2002 Total	3-yr Average
			9-May	4-Jun	9-Jul	6-Aug	18-Sep	29-Oct		
NM9D11A-PAR	4.44*	9.85*	0.75	1.35*	2.15*	1.81	1.53	1.04	8.62*	7.64**
NC+605	4.48**	9.47*	0.93	1.34*	2.09	1.73*	1.47*	1.04	8.59*	7.51*
Wilson	4.27*	8.96*	0.89	1.38*	2.31*	1.75*	1.73**	1.15	9.21**	7.48*
Rio Grande	4.11*	9.92**	0.87	1.30*	2.07	1.76*	1.46*	0.88	8.33*	7.46*
Magna 601	3.80*	9.54*	0.65	1.38*	2.31*	1.95**	1.57*	1.01	8.86*	7.40*
5681	4.25*	9.67*	0.83	1.32*	2.27*	1.48*	1.37	0.83	8.09*	7.34*
Archer II	3.95*	9.72*	0.77	1.25*	2.14*	1.73*	1.52*	0.84	8.25*	7.30*
ZX9362	4.06*	9.36*	0.77	1.25*	2.11*	1.76*	1.44*	0.88	8.20*	7.21*
Dona Ana	3.34	9.04*	0.62	1.18*	1.74	1.67*	1.33	2.02**	8.56*	6.98*
WL 327	3.30	9.35*	0.80	1.41*	2.07	1.61*	1.30	0.78	7.97*	6.87*
Cimarron 3i	3.09	8.91*	1.37**	1.49**	2.08	1.75*	1.15	0.71	8.53*	6.85*
Abilene + Z	3.33	9.28*	0.76	1.33*	2.06	1.58*	1.33	0.79	7.83	6.81*
Ram	3.49	8.95*	0.88	1.37*	1.98	1.54*	1.31	0.88	7.95*	6.80*
WL 442	4.01*	8.39	0.54	1.05*	2.04	1.68*	1.59*	1.09	7.98*	6.79*
54Q53	2.95	8.99*	0.77	1.37*	2.06	1.86*	1.37	0.82	8.23*	6.72*
Sutter	3.01	9.14*	0.73	1.29*	1.99	1.75*	1.42*	0.79	7.96*	6.71*
NM Common	3.22	9.16*	0.80	1.26*	1.95	1.61*	1.30	0.77	7.68	6.68*
Dagger + EV	3.03	9.14*	0.67	1.26*	2.12*	1.71*	1.16	0.73	7.65	6.60
Garst 6420	2.72	9.10*	0.86	1.22*	2.04	1.71*	1.45*	0.69	7.98*	6.60
SD-Common	2.40	9.04*	1.01	1.34*	2.05	1.81*	1.40	0.50	8.12*	6.52
DK142	2.81	8.74	0.72	1.10*	2.43**	1.66*	1.24	0.65	7.79	6.45
Magnum V	2.78	8.92*	0.85	1.20*	1.95	1.57*	1.22	0.55	7.34	6.35
Garst 6550	2.75	8.80	0.50	1.27*	2.05	1.56*	1.31	0.75	7.44	6.33
ABT400SCL	2.88	8.54	0.76	1.21*	1.87	1.52*	1.20	0.68	7.24	6.22
ABT 350	2.92	8.17	0.61	1.19*	1.99	1.71*	1.28	0.59	7.36	6.15
GH 766	2.75	8.38	0.50	1.04*	1.99	1.53*	1.25	0.62	6.93	6.02
GH 750	2.29	8.57	0.68	1.10*	1.94	1.51*	1.27	0.62	7.12	5.99
PGI 4372	1.70	7.98	0.85	1.01*	1.70	1.46*	1.11	0.49	6.62	5.43
Mean	3.29	9.04	0.78	1.26	2.06	1.67	1.36	0.83	7.94	6.76
LSD (0.05)	1.04	1.04	0.23	Ns	0.34	Ns	0.32	0.62	1.29	0.98
CV%	22.39	8.17	21.25	16.06	11.60	13.84	16.90	53.61	11.51	17.40

¹Yields in 2002 for this test are exceptionally high compared to other tests managed similarly and located within 1/8 mile (see table 8). Field drainage and soil water-holding characteristics in around this test are apparently very conducive to alfalfa production compared to other areas of this research facility.

²Yield data from previous years may be different than that presented in other publications due to a difference in analysis methods. 2000 Harvest dates: 21-Jun, 20-Jul, 17-Aug, 11-Sep, and 2-Nov.

2001 Harvest dates: 10-May, 11-Jun, 9-Jul, 7-aug, 10-Sep, and 29-Oct.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

LSD (0.05) stands for the Least Significant Difference at the 5% level. If the difference between two numbers within a column is equal to or greater than the LSD, it is 95% certain that they are truly different.

Ns means that there were no significant differences between varieties within that column at the 5% level.

Table 8. Dry matter yields (tons/acre) of alfalfa varieties sown August 30, 2001, at NMSU's Agricultural Science Center at Tucumcari and scheduled to be furrow-irrigated once per cutting but only on May 15 in 2001 due to water shortages.

Variety	Yield (dry tons/acre)			2002 Total
	5-Jun	9-Jul	6-Aug	
Magna 601	1.42	1.63**	0.87*	3.91**
HybriGreen 41	1.72**	1.59*	0.59	3.90*
Wilson	1.53*	1.50*	0.82*	3.85*
NM9D11A-PAR	1.36	1.54*	0.90**	3.80*
Archer	1.38	1.62*	0.77*	3.78*
RSC-681	1.70*	1.44*	0.62	3.76*
NM Common	1.37	1.45*	0.87*	3.69*
NC+Jade II	1.43	1.56*	0.70*	3.69*
Express	1.33	1.56*	0.78*	3.67*
Tango	1.34	1.51*	0.77*	3.61*
HybriForce 400	1.43	1.46*	0.72*	3.61*
African Common	1.20	1.48*	0.88*	3.56*
Haygrazer	1.54*	1.43*	0.58	3.54*
Dona Ana	1.30	1.44*	0.72*	3.46*
Forecast 1001	1.41	1.30*	0.73*	3.43*
AmeriStand 403T	1.23	1.49*	0.65	3.37*
WR9801	1.47*	1.31*	0.50	3.29*
Select	1.32	1.37*	0.60	3.29*
Mean	1.42	1.48	0.73	3.62
LSD (0.05)	0.29	Ns	0.23	Ns
CV%	12.22	10.05	19.40	7.97

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

LSD (0.05) stands for the Least Significant Difference at the 5% level. If the difference between two numbers within a column is equal to or greater than the LSD, it is 95% certain that they are truly different.

Ns means that there were no significant differences between varieties within that column at the 5% level.

Table 9. Dry matter yields (tons/acre) of flood-irrigated alfalfa varieties sown September 1, 1999, at NMSU's Agricultural Science Center at Los Lunas and flood-irrigated twice per cutting.

Variety	2000 Total	2001 Total	2002 Harvests					2002 Total	3-yr Average
			16-May	2-Jun	25-Jul	2-Sep	24-Oct		
CW 55112	11.41**	11.62*	2.98*	2.97*	2.93*	2.49**	1.32*	12.75**	11.90**
Artesian Sunrise	11.25*	11.74**	3.11**	2.91*	2.67*	2.48*	1.32*	12.45*	11.82*
CW 5567	11.00*	11.58*	3.11**	3.06**	2.64*	2.45*	1.19*	12.58*	11.68*
CW 6699	10.14*	11.18*	3.07*	2.77	2.96**	2.24*	1.36**	12.33*	11.22
Mean	10.95	11.53	3.07	2.93	2.80	2.42	1.30	12.53	11.66
LSD (0.05)	Ns	Ns	Ns	0.16	Ns	Ns	Ns	Ns	0.47
CV%	5.70	4.85	7.31	3.72	10.36	8.95	16.85	4.23	4.84

¹Yield data from previous years may be different than that presented in other publications due to a difference in analysis methods.

2000 Harvest dates: 3-May, 14-Jun, 20-Jul, and 21-Aug.

2001 Harvest dates: 10-May, 11-Jun, 9-Jul, 7-Aug, 10-Sep, and 29-Oct.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

LSD (0.05) stands for the Least Significant Difference at the 5% level. If the difference between two numbers within a column is equal to or greater than the LSD, it is 95% certain that they are truly different.

Ns means that there were no significant differences between varieties within that column at the 5% level.

Table 10. Dry matter yields (tons/acre) of alfalfa varieties sown August 28, 2001, at NMSU's Agricultural Science Center at Los Lunas and flood-irrigated twice per cutting.

Variety	2002 Harvests					2002 Total
	16-May	2-Jun	25-Jul	2-Sep	21-Oct	
WR9801	1.03*	1.19*	1.31*	1.35**	0.73*	5.59**
HybriForce 400	0.92	1.19*	1.33*	1.31*	0.80*	5.55*
Cimarron 3i	1.15**	1.13*	1.31*	1.21*	0.71*	5.51*
Magna 601	0.80	1.23*	1.36**	1.25*	0.84*	5.47*
HybriGreen 41	0.96	1.19*	1.28*	1.24*	0.77*	5.44*
Rio	0.76	1.29**	1.27*	1.17*	0.84*	5.33*
Aspire	0.69	1.16*	1.27*	1.29*	0.85**	5.26*
Archer	0.77	1.16*	1.27*	1.25*	0.81*	5.26*
NM9D11A-PAR	0.82	1.12*	1.21*	1.15*	0.79*	5.09*
NM Common	0.63	1.10*	1.25*	1.10*	0.70*	4.78*
Mean	0.85	1.18	1.29	1.23	0.78	5.33
LSD (0.05)	0.15	Ns	Ns	Ns	Ns	Ns
CV%	11.97	8.81	8.11	10.63	12.50	7.80

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

LSD (0.05) stands for the Least Significant Difference at the 5% level. If the difference between two numbers within a column is equal to or greater than the LSD, it is 95% certain that they are truly different.

Ns means that there were no significant differences between varieties within that column at the 5% level.

Table 11. Dry matter yields (tons/acre) of alfalfa varieties sown 7 September 2001, at the NMSU's Sustainable Agriculture Science Center at Alcalde and furrow-irrigated once per cutting.¹

Variety	2002 Harvests		2002 Total
	18-Jul	27-Aug	
Tango	1.65*	1.40*	3.04**
BPR379	1.54*	1.46**	3.00*
NM Common	1.69**	1.27*	2.96*
Archer	1.60*	1.27*	2.87*
HybriGreen 41	1.52*	1.35*	2.86*
ZG0152A	1.53*	1.31*	2.85*
HybriForce 400	1.53*	1.31*	2.84*
AmeriGraze 401+Z	1.46*	1.35*	2.81*
SS120	1.46*	1.34*	2.80*
ZG0160A	1.45*	1.31*	2.75*
Deepkrown	1.45*	1.29*	2.74*
MagnaGraze	1.51*	1.22	2.72*
Wilson	1.47*	1.22	2.69*
DU-201	1.36*	1.32*	2.68*
Spredor-3	1.54*	1.13	2.67*
African Common	1.47*	1.20	2.66*
Haygrazer	1.42*	1.22	2.64*
AmeriStand 403T	1.40*	1.20	2.61*
AlfaGraze	1.44*	1.11	2.55*
CO Common	1.28*	1.06	2.35*
Mean	1.49	1.27	2.76
LSD (0.05)	Ns	0.20	Ns
CV%	21.97	15.95	15.86

¹The test was planted to compare varieties with and without continuous winter grazing pressure. First cutting yields were removed on and discarded on 18-Jun because of heavy weed infestation. Grazing by 4 mixed breed heifers weighing approximately 500 lb was initiated on 26 November and will continue until mid-May 2003. There was no difference between yields of grazed and ungrazed treatments in 2002 and no grazing treatment X variety interaction.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

LSD (0.05) stands for the Least Significant Difference at the 5% level. If the difference between two numbers within a column is equal to or greater than the LSD, it is 95% certain that they are truly different.

Ns, not significantly different based on the 5% LSD.

Table 12. Dry matter yields (tons/acre) of alfalfa varieties sown August 13, 1999, at NMSU's Agricultural Science Center at Farmington and sprinkler-irrigated three times per week.

Variety	2000 Total	2001 Total	2002 Harvests				2002 Total	3-yr Aver- age
			29-May	3-Jul	8-Aug	24-Sep		
DK142	8.18*	9.59*	3.41*	2.60*	2.01*	1.92*	9.93*	9.23**
Affinity+Z	7.90*	9.25*	4.08**	2.61*	1.95*	1.76*	10.39**	9.18*
Millennia	7.80*	9.88**	3.30*	2.43*	1.96*	2.03**	9.72*	9.13*
Geneva	8.23*	9.41*	3.52*	2.50*	1.95*	1.77*	9.74*	9.13*
Select	8.35**	9.43*	3.28*	2.13*	1.89*	1.75*	9.05*	8.94*
Archer II	7.30*	9.66*	3.43*	2.43*	2.00*	1.82*	9.68*	8.88*
ZX9351	7.45*	9.06*	3.63*	2.47*	2.02*	1.93*	10.05*	8.85*
Focus	7.93*	8.92*	3.42*	2.34*	1.98*	1.85*	9.59*	8.81*
Somerset	7.25*	8.96*	3.56*	2.62**	2.13**	1.74*	10.05*	8.76*
Mean	7.82	9.35	3.51	2.46	1.99	1.84	9.80	8.99
LSD (0.05)	Ns	Ns	Ns	Ns	Ns	Ns	Ns	Ns
CV%	12.01	5.10	12.27	15.28	10.52	11.14	9.53	9.01

¹Yield data from previous years may be different than that presented in other publications due to a difference in analysis methods.

2000 Harvest dates: 31-May, 6-Jul, 8-Aug, and 21-Sep.

2001 Harvest dates: 1-Jun, 5-Jul, 8-Aug, and 20-Sep.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

LSD (0.05) stands for the Least Significant Difference at the 5% level. If the difference between two numbers within a column is equal to or greater than the LSD, it is 95% certain that they are truly different.

Ns means that there were no significant differences between varieties within that column at the 5% level.

Table 13. Dry matter yields (tons/acre) of alfalfa varieties sown August 23, 2001, at NMSU's Agricultural Science Center at Farmington and sprinkler-irrigated three times per week.

Variety	Yield (dry tons/acre)				2002 Total
	29-May	3-Jul	6-Aug	24-Sep	
HybriGreen 41	2.82*	2.28*	1.50*	1.59**	8.18**
Archer II	2.91*	2.14*	1.34	1.56*	7.95*
54V54	2.92*	1.98*	1.34	1.36*	7.59*
HybriForce 400	2.56*	2.12*	1.55**	1.35*	7.59*
DKA42-15	2.50*	2.37*	1.34	1.38*	7.58*
Forecast 1001	2.75*	1.98*	1.47*	1.34*	7.55*
Legend	2.61*	2.08*	1.41*	1.43*	7.53*
WL 342	2.30	2.57**	1.31	1.34*	7.53*
Dona Ana	2.52*	2.17*	1.40*	1.41*	7.50
Champ	2.44	2.16*	1.40*	1.44*	7.45
Focus	2.70*	1.96*	1.46*	1.32*	7.44
Megaton 3.5	2.19	2.37*	1.49*	1.39*	7.43
5-Star	2.56*	2.14*	1.42*	1.29*	7.40
WL 327	2.45	2.14*	1.29	1.33*	7.20
Geneva	2.28	2.18*	1.38	1.31*	7.15
Magnum V	2.14	2.22*	1.38	1.29*	7.04
NM9D11A-PAR	1.92	2.24*	1.30	1.44*	6.89
Delta 526	2.02	2.13*	1.26	1.24*	6.64
Mean	2.48	2.18	1.39	1.38	7.42
LSD (0.05)	0.45	Ns	0.16	Ns	0.68
CV%	12.91	14.63	7.87	12.54	6.46

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

LSD (0.05) stands for the Least Significant Difference at the 5% level. If the difference between two numbers within a column is equal to or greater than the LSD, it is 95% certain that they are truly different.

Ns means that there were no significant differences between varieties within that column at the 5% level.

Table 14. Characteristics and performance of alfalfa varieties across years and tests in New Mexico.		Varietal Characteristics ¹										Las Cruces			Artesia				Tucumcari				Los Lunas				Alcalde		Farmington								
		WS	FD	Pest resistance								1999 ²			1999			1999 ³			01	1999			01	2000 ⁴			1999			01					
				BW	PRR	FW	AN	SAA	PA	BAA	00 ⁵	01	02	00	01	02	00	01	02	02	00	01	02	02	00	01	02	02	00	01	02	02					
Variety	Proprietor																																				
AmeriStand 801S	America's Alfalfa	n/r	8	R	R	HR	R	HR	MR	HR				*	*	*																					
AmeriStand 802	America's Alfalfa	n/r	8	?	HR	?	R	?	?	HR				*	*	**				*																	
C/W 5875	CalWest Seeds	n/r	8	HR	HR	HR	HR	HR	HR	HR	*	*	*	*	*	*																					
C/W 68115	CalWest Seeds	n/r	8	HR	HR	HR	HR	HR	HR	HR	*	*	**	*	*	*																					
C/W 78122	CalWest Seeds	n/r	8	HR	HR	HR	HR	HR	?	HR	**	*	*	*	*	*																					
CutMor	Union Seed Co.	6.0	8	LR	HR	HR	R	HR	HR	HR	*			*	*	*																					
DK180ML	Monsanto	6.0	8	MR	HR	HR	HR	HR	HR	HR				*	*	*																					
DS8181	Kelly Green Seed	n/r	8	?	?	?	?	?	?	?																											**
Duran 2-2	Great Plains Research	n/r	8	R	HR	HR	HR	HR	HR	R																										*	
GT 13-R Plus	America's Alfalfa	n/r	8	R	R	HR	?	R	MR	LR	*	*	*																								
Magna 801FQ	Dairyland Seed Co.	n/r	8	R	HR	HR	MR	HR	R	R				*	*	*																					
Mesa	PGI Alfalfa	5.4	8	MR	R	HR	?	HR	HR	R	*	*	*																								
Pershing	PGI Alfalfa	5.3	8	HR	HR	HR	LR	HR	R	R	*																										
PGI 8000	PGI Alfalfa	5.5	8	?	?	?	?	?	?	?	*	*																									
Rio Grande	Great Plains Research	n/r	8	MR	HR	HR	HR	HR	HR	HR	*	*					*	**	*		*	**	*		*	**	*										
Signal 8000	Helena Chemical	?	8	?	?	?	?	?	?	?	*	**					*	*	*		*	*	*														
WL525HQ	WL Research	n/r	8	MR	HR	MR	?	HR	HR	HR				*	*	*	*	*	*		*	*	*														
ZG9891	ABI Alfalfa	n/r	8	?	R	HR	HR	?	?	R		*																									
ZX9889B	ABI Alfalfa	n/r	8	?	R	?	R	?	?	R	*	*																									
Magna 901	Dairyland Seed Co.	n/r	9	MR	HR	HR	R	HR	R	HR				*	*	*	*	*	*	*		*	*	*													
Salado	America's Alfalfa	n/r	9	?	LR	R	LR	R	MR	HR							*	*	*		*	*	*														
WL612	WL Research	n/r	9	?	HR	HR	LR	HR	HR	HR	*	*																									
WL625HQ	WL Research	n/r	9	MR	HR	HR	LR	HR	HR	HR																											
ZS9992	ABI Alfalfa	n/r	9	?	R	R	HR	?	?	?																											
ZX9393	ABI Alfalfa	n/r	9	MR	R	HR	R	HR	R	HR				*	*	*																					
ZX9894	ABI Alfalfa	n/r	9	?	R	HR	R	?	HR	R																											
ZS0000	ABI Alfalfa	n/r	#	?	R	HR	?	R	?	R																											

¹WS=Winter Survival (1=No injury, 6=Dead plants; n/r=not rated, generally for FD>6; be cautious when using this data because most of the testing is done in the northern U.S. and might underestimate winter survivability in New Mexico), FD=Fall Dormancy (2=Vernal, 3=Ranger, 4=Saranac, 5=Archer, 6=ABI 700, 7=Dona Ana, 8=Maricopa, 9-CUF 101), Bw=Bacterial wilt, PRR=Phytophthora root rot, Fw=Fusarium wilt, An=Anthracnose, SAA=Spotted alfalfa aphid, PA=Pea aphid, BAA=Blue alfalfa aphid (S=Susceptible, LR=Low Resistance, MR=Moderate Resistance, R=Resistant, HR=High Resistance).

²Establishment year.

³Test was drip irrigated.

⁴Two tests were sown in 2001 at Alcalde, one to be grazed from November to May beginning in fall 2002. Both will be managed for hay production during the growing season. There were no differences between n tests in 2002 and not variety X test interactions.

⁵Harvest Year.

Shaded boxes indicate that the variety was not in the test.

**Highest yielding variety in the test for that year.

*Not significantly different from the highest yielding variety in the test for that year.

L.M. Lauriault, I.M. Ray, D.A. McWilliams, L.M. English, R.P. Flynn, S.J. Guldán, M.W. Murray, and M.K. O'Neill
 New Mexico State University College of Agriculture and Home Economics Agricultural Experiment Station and Cooperative Extension Service

Table 15. New Mexico State University Cooperative Extension Service publications related to alfalfa management.

Number	Title	Online ?
A-107	Managing saline soils	
A-113	Selection of fertilizers	Y
A-114	Test your soil	Y
A-122	Soil test interpretations	Y
A-123	Sampling for plant tissue analysis	
A-128	Fertilizer guide for New Mexico	Y
A-128	Nitrogen fixation by legumes	Y
A-130	Inoculation of legumes	Y
A-131	Certified seed	Y
A-133	Calculating fertilizer costs	Y
A-134	Selecting synthetic fertilizers in New Mexico	Y
A-18	Micronutrient fertility guide	
A-216	Know what is in a bag of seed	Y
A-309	Alfalfa weevil and clover leaf weevil	
A-316	Structure of a hay bale	
A-317	Alfalfa fertilization in New Mexico	
A-318	Reducing alfalfa harvest losses	Y
A-325	Managing weeds in alfalfa	Y
A-327	Introduction to hay testing	Y
A-328	Sampling guidelines for hay testing	Y
A-329	Variations in hay grading	Y
A-330	Alfalfa growth stages	Y
A-331	Alfalfa quality definitions	Y
B-115	Balancing forage supply and demand	Y
CR-536	Blister beetles in alfalfa	Y
CR-581	Drought strategies for alfalfa	Y
HB-11	Suggestions for managing insects in alfalfa and clover 1996	
W-01	Submitting plants for plant tissue analysis	
W-13	Alfalfa disease control	

These publications are available from your county office of the NMSU Cooperative Extension Service or online from the Internet at <http://www.cahe.nmsu.edu/pubs/>