



The 2017
New Mexico
Alfalfa Variety
Test Report



Agricultural Experiment Station
**College of Agricultural, Consumer
and Environmental Sciences**



The 2017 New Mexico Alfalfa Variety Test Report

Leonard Lauriault, Ian Ray, Chris Pierce, Owen Burney, Robert Flynn, Mark Marsalis, Mick O'Neill, Ashley Cunningham, Charles Havlik, and Margaret West¹

Introduction

In 2017, 190,000 acres of alfalfa (*Medicago sativa*) were in production in New Mexico, which was unchanged from 2016. Hay yields were estimated at 950,000 tons reflecting a 2% increase in yield. At a January through November 2017 average of \$177/ton (up from \$170/ton in 2016), estimated gross returns from alfalfa hay produced in 2017 will total just over \$168 million, a \$10 million increase over 2016, but still below the \$188 million received in 2015. Besides its value for hay, alfalfa also is the legume of choice in irrigated perennial pastures. Whether used as pasture or hay, the value of alfalfa to New Mexico is greatly magnified by its contribution to livestock production and receipts from the sale of meat, milk, and other products generated by livestock enterprises.

Choosing a good alfalfa variety is a key step in establishing a highly productive stand of alfalfa, whether for hay or pasture. Differences between the highest- and lowest-yielding varieties in irrigated tests included in this report ranged from 0.90 to 2.05 tons per acre in 2017. If sold as hay, this translates to a potential difference in returns of \$159 to \$363 per acre due to variety, or an increase of at least \$30 million for the industry in 2017 alone.

This report, which is a collaborative effort of New Mexico State University scientists at agricultural science centers throughout the state, provides yield data for alfalfa varieties included in yield trials in New Mexico. While consistently high yields compared to other varieties over a number of years and locations within a region is the best indication of varietal adaptation and persistence, other factors should be considered in the variety selection process (see NMSU's Cooperative Extension Service Circular 654, *Selecting alfalfa varieties for New Mexico*). In addition to fall dormancy and winter hardiness, high levels of pest resistance are critical to protecting an alfalfa stand for long-term production. Alfalfa grown in New Mexico should have at least a resistant (R) rating for bacterial wilt, Fusarium wilt, anthracnose, Phytophthora root rot, spotted alfalfa aphid, blue alfalfa aphid, pea aphid, stem nematode, and southern rootknot nematode. Seed quality also should be high. Selecting an alfalfa variety based on seed cost is a gamble producers often lose. To be assured of achieving a long-lasting, highly productive stand, buy either certified or Plant Variety Protected (PVP) seed, which guarantees the genetics and performance. The best choice of seed of any variety is one that was treated with a fungicide and nitrogen-fixing bacteria before it was bagged.

Description of Tests

Replicated alfalfa variety tests included in this report were conducted under research controls at NMSU's Agricultural Science Centers at Las Cruces (2014 normal, drought, and early termination irrigation studies), [2014 (Artesia late spring planted) and 2016], Tucumcari (2015 irrigated with treated municipal wastewater), Los Lunas (2016), and Farmington

(2014). Weather data for 2017 and the long-term averages from all locations are presented in table 1.

Yield data (on a dry matter basis) are presented in tables 2-9. Varieties are listed in order from highest to lowest average annual production. Yields are given by cutting for 2017 and by year for each production year. Statistical analyses were performed on all alfalfa yield data (including experimental entries) to determine if the apparent differences are truly due to variety or just to chance. The variety with the highest numerical yield in each column is marked with two asterisks (**), and those varieties not significantly different from that variety are marked with one asterisk (*). Those are the varieties from which to make an initial selection. Otherwise, to determine if two varieties are truly different, compare the difference between the two varieties to the Least Significant Difference (LSD) at the bottom of the column. If the difference is equal to or greater than the LSD, the varieties are truly different in yield when grown under the conditions at a given location. If NS is given for the LSD, there was no statistical difference between the highest and lowest yielding varieties. The Coefficient of Variation (CV), which is a measure of the variability of the data, is included for each column of means. Low variability (<20 percent) is desirable, and increased variability within a study results in higher CVs and larger LSDs. There might be a difference between previously published data and the data given in this publication for the same tests because of differences in the programs used for statistical analysis.

Table 10 summarizes information about proprietors, Roundup Ready genetics, fall dormancy, winter survival (measured in the northern United States), pest resistance, and yield performance across years and locations for all varieties currently included in NMSU's alfalfa variety testing program. Varieties are listed alphabetically by fall dormancy category. As in the data tables, the variety with the highest numerical yield in each column is marked with two asterisks (**), and those varieties not significantly different from that variety are marked with one asterisk (*). Remember good performance across several years and locations is the best indicator of broad adaptation, pest resistance, and persistence.

Seed labeled "common," "variety not stated," or "variety unknown", particularly that from other states, is of unknown genetic background and may or may not have the necessary disease or insect resistance. New Mexico Common and African Common seed used in all tests throughout the state has come from the same supplier and seed fields in New Mexico. Seed purchased from other dealers may or may not be of the same quality and performance.

Summary

Consistent production of high alfalfa yields is the result of selecting good varieties and implementing good management techniques. Soil fertility should be maintained at recommended levels based on soil tests, irrigation should be properly applied, weeds and insects should be controlled using appropriate cultural and/or chemical methods, and harvest man-

¹Forage Crop Management Scientist and Superintendent, NMSU Agricultural Science Center at Tucumcari; Alfalfa Breeder, NMSU, Las Cruces; Forage Research Scientist, NMSU, Las Cruces; Superintendent, NMSU John T. Harrington Forestry Research Center at Mora; Extension Agronomist, NMSU Agricultural Science Center at Artesia; Extension Forage Specialist and Superintendent, NMSU Agricultural Science Center at Los Lunas; Agronomist, NMSU Agricultural Science Center at Farmington; Research Assistant, NMSU Agricultural Science Center at Tucumcari; Senior Research Assistant, NMSU Agricultural Science Center at Los Lunas, and Agricultural Research Scientist, NMSU Agricultural Science Center at Farmington, respectively.

agement should allow sufficient time to restock root energy prior to winter. For dormant (FD 1 to 3) and semidormant (FD 4 to 6) varieties, a 6-week rest period before a dormancy-inducing freeze (27°F) is recommended to allow plants to replenish root reserves for winter survival and initiate spring growth, after which harvesting might be done either mechanically or by grazing. Non-dormant (FD 7 to 9) varieties also might benefit from this rest period. Removing fall growth is beneficial to reducing weevil populations the following year as eggs are laid in and overwinter in stems. Harvesting established stands at early bloom would result in 3 to 5 cuttings per year before initiation of the rest period in most areas of New Mexico. More dormant varieties might not produce yields that can be baled during the rest period; however, these can still be grazed. For further information about alfalfa management, refer to the other NMSU Agricultural Experiment Station and Cooperative Extension Service publications listed in table 11.

Acknowledgements

The authors express appreciation to the following for their significant contribution to The New Mexico Alfalfa Variety Testing Program by helping with planting, maintaining, harvesting, or other data collection, and data entry: Dallen Begay, Nathan Begay, Jason Box, Servando Bustillos, Patty Cooksey, Josh Foster, Jared Jennings, Shane Jennings, Jonah Joe, Ruben Pacheco, Tom Place, Dennis Price, Mario Segura, Anthony Williams, and the staff at University Communications who make publications such as this possible. Salaries and research support were provided by state and federal funds appropriated to the New Mexico Agricultural Experiment Station. Federal funds included those appropriated through the Hatch Act of 1887.

Table 1. Temperature and precipitation data for 2017 and the long-term averages for the New Mexico Alfalfa Variety Test locations.

Location Elevation Latitude	Las Cruces 3832 ft. 32° 12' N				Artesia 3366 ft. 32° 45' N				Tucumcari 4091 ft. 35° 12' N				Los Lunas 4840 ft. 34° 46' N				Farmington 5640 ft. 36° 41' N			
	Temp. (°F)		Precip. (in.)		Temp. (°F)		Precip. (in.)		Temp. (°F)		Precip. (in.)		Temp. (°F)		Precip. (in.)		Temp. (°F)		Precip. (in.)	
Month	2017	Avg.	2017	Avg.	2017	Avg.	2017	Avg.	2017	Avg.	2017	Avg.	2017	Avg.	2017	Avg.	2017	Avg.	2017	Avg.
Nov-16	52	49	0.02	0.53	52	49	0.73	0.53	51	47	1.39	0.71	46	44	1.46	0.46	45	41	1.50	0.65
Dec-16	45	41	0.73	0.68	42	41	0.76	0.51	39	39	0.37	0.67	36	35	0.70	0.52	34	31	0.97	0.47
Jan-17	45	40	1.50	0.56	44	40	0.89	0.39	38	38	1.02	0.42	37	35	1.43	0.36	33	30	0.71	0.51
Feb-17	51	45	0.36	0.37	50	45	0.41	0.42	48	42	0.17	0.49	44	40	0.94	0.41	41	36	0.93	0.48
Mar-17	60	52	0.00	0.22	58	52	0.02	0.43	56	49	2.16	0.76	52	47	0.16	0.50	49	43	0.39	0.64
Apr-17	65	60	0.07	0.21	63	60	1.09	0.62	58	58	2.73	1.14	55	55	0.87	0.46	50	51	1.51	0.58
May-17	70	69	0.13	0.29	67	69	0.30	1.20	64	66	1.82	1.94	63	63	0.42	0.45	58	60	1.14	0.53
Jun-17	80	78	0.07	0.72	80	78	1.83	1.40	77	76	0.98	1.98	76	72	0.05	0.61	73	70	0.02	0.26
Jul-17	79	80	4.32	1.36	81	80	1.49	1.76	82	79	1.58	2.75	79	77	0.60	1.25	77	76	2.51	0.81
Aug-17	77	78	1.95	2.29	78	78	3.15	1.67	75	77	6.48	2.78	74	75	3.12	1.70	74	74	0.11	1.05
Sep-17	73	71	0.47	1.38	73	71	1.92	1.81	70	71	2.65	1.62	68	67	1.71	1.17	67	66	1.57	1.10
Oct-17	63	61	0.28	0.91	62	61	0.43	1.16	60	60	3.62	1.31	55	56	1.11	1.04	55	54	0.08	0.97
Annual	63	60	9.90	9.52	62	60	13.02	11.90	60	58	24.97	16.57	57	56	12.57	8.93	55	53	11.44	8.05

Table 2. Dry matter yields (tons/acre) of alfalfa varieties sown October 7, 2014, at NMSU's Leyendecker Plant Science Research Center at Las Cruces and flood-irrigated twice per cutting to apply 58.16 inches (normal irrigation)†.

Variety Name	2015 Total	2016 Total	2017 Harvests						2017 Total	3-Yr Average
			27-Apr	24-May	26-Jun	9-Aug	13-Sep	17-Oct		
AmeriStand 803T	10.07**	8.22**	1.65*	1.05*	1.11*	1.58*	0.96*	0.87**	7.24*	8.36**
WL 656HQ	8.98	8.01*	1.61	1.15**	1.20*	1.64*	0.92*	0.85*	7.38*	8.33*
NM14BMHS1	9.93*	7.80*	1.78*	0.94	0.82	1.49*	0.77	0.63	6.37*	8.21*
NM14BMC0	9.85*	7.59*	1.86*	0.97	0.97	1.51*	0.76	0.70	6.81*	8.10*
NM14GTAF07235	9.42*	7.71*	1.59	1.06*	0.91	1.54*	0.87*	0.82*	6.80*	8.04*
4N900	9.00*	7.65*	1.65*	1.04*	1.05*	1.63*	0.97*	0.87**	7.20*	7.88*
NM14BMHS3	9.42*	7.37*	1.64	0.87	0.83	1.50*	0.73	0.63	6.18	7.84*
FSG903	9.33*	7.67*	1.55	0.97	0.98	1.49*	0.82	0.81*	6.65*	7.77*
NM14MaiHS3	9.85*	7.49*	1.81*	0.86	0.78	1.43*	0.69	0.69	6.28	7.76*
NuMex Bill Melton	8.53	7.46*	1.78*	1.01	1.26**	1.67*	0.92*	0.77*	7.41**	7.74*
Malone	8.65	7.28*	1.44	1.08*	1.10*	1.70**	1.05**	0.86*	7.21*	7.72*
NM14MLLS2	8.85	7.43*	1.70*	0.83	0.91	1.52*	0.79	0.65	6.39*	7.55*
NM1407227	8.75	7.39*	1.87**	0.84	0.90	1.40*	0.69	0.59	6.32*	7.55*
NM14BM1008251	8.71	7.20*	1.60	0.97	1.03*	1.59*	0.93*	0.79*	6.91*	7.54*
AFX149092	8.80	7.19*	1.46	1.02*	0.92	1.42*	0.76	0.75*	6.30	7.47*
NM14ALWLHQ	9.06	7.18*	1.58	0.93	0.88	1.38*	0.72	0.65	6.16	7.43*
Sandpiper	8.52	7.01*	1.61	0.89	0.92	1.53*	0.80	0.67	6.45*	7.42*
Cisco II	9.42*	6.90*	1.55	0.82	0.74	1.37*	0.73	0.65	5.87	7.38*
WL 440HQ	8.99	6.97*	1.60	0.88	0.78	1.32*	0.65	0.58	5.83	7.35*
NM14BMHR2	9.15	7.05*	1.75*	0.85	0.88	1.51*	0.72	0.62	6.27	7.33*
AFX148091	8.60	6.92*	1.40	0.90	0.86	1.36*	0.70	0.65	5.82	7.17*
Hi-Gest 660	8.88	6.98*	1.43	0.87	0.84	1.30*	0.70	0.67	5.82	7.13*
Wilson	8.88	6.68*	1.53	0.89	0.82	1.43*	0.87*	0.79*	6.31	7.12*
57Q53	7.90	6.56*	1.63	0.88	0.87	1.33*	0.58	0.56	5.84	6.84*
Mean	9.06	7.32	1.63	0.94	0.93	1.49	0.80	0.71	6.49	7.63
LSD (0.05)	0.81	NS	0.23	0.14	0.26	NS	0.22	0.14	1.10	NS
CV%	6.80	8.82	9.84	10.40	19.44	12.44	19.76	13.91	11.97	14.83

†Data were analyzed using analysis of covariance where check plots of NuMex Bill Melton were used as the covariate.

2015 Harvest dates: 28-Apr, 12-Jun, 16-Jul, 12-Aug, 9-Sep, and 6-Nov.

2016 Harvest dates: 27-Apr, 1-Jun, 28-Jun, 28-Jul, 14-Sep, and 26-Oct.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 3. Dry matter yields (tons/acre) of alfalfa varieties sown October 7, 2014, at NMSU's Leyendecker Plant Science Research Center at Las Cruces and flood-irrigated once per cutting to apply 41.31 inches (drought irrigation)†.

Variety Name	2015 Total	2016 Total	2017 Harvests						2017 Total	3-Yr Average
			1-May	26-May	21-Jun	8-Aug	12-Sep	18-Oct		
AmeriStand 803T	7.07*	3.83*	1.00*	0.48*	0.20*	1.04*	0.60*	0.51*	3.76*	4.85**
Wilson	6.45*	3.75*	0.96*	0.41*	0.22*	1.05*	0.66**	0.53**	3.86*	4.81*
NuMex Bill Melton	6.27*	3.79*	1.10**	0.53*	0.28**	1.07*	0.65*	0.50*	4.13**	4.74*
FSG903	6.69*	3.84**	0.98*	0.41*	0.19*	0.97*	0.49*	0.41*	3.51*	4.73*
NM14BMHS1	7.29*	3.51*	1.02*	0.31	0.10*	0.92	0.43*	0.33	3.10	4.67*
NM14BM1008251	6.95*	3.52*	0.95	0.39	0.09*	0.98*	0.46*	0.39*	3.27*	4.59*
AFX149092	6.80*	3.40*	0.91	0.40	0.09*	0.87	0.37	0.37*	3.06	4.55*
WL 656HQ	5.86	3.73*	0.95	0.54*	0.21*	1.05*	0.64*	0.52*	3.90*	4.54*
4N900	6.78*	3.56*	0.96*	0.56**	0.19*	0.96*	0.45*	0.42*	3.51*	4.49*
Malone	5.77	3.64*	0.80	0.51*	0.26*	1.13**	0.62*	0.48*	3.85*	4.49*
NM1407227	6.85*	3.28*	1.09*	0.30	0.10*	0.99*	0.44*	0.33	3.28*	4.42*
NM14BMHS3	7.44**	3.32*	0.95	0.26	0.10*	0.75	0.24	0.20	2.54	4.40*
NM14MaHS3	6.67*	3.50*	1.08*	0.31	0.10*	0.90	0.35	0.30	3.00	4.38*
NM14GTAF07235	6.44*	3.50*	0.90	0.42*	0.15*	0.87	0.44*	0.36*	3.15	4.32*
NM14ALWLHQ	6.31*	3.33*	0.98*	0.36	0.09*	0.90	0.44*	0.35*	3.15	4.27*
NM14BMC0	6.39*	3.40*	1.08*	0.39	0.11*	1.00*	0.44*	0.35*	3.30*	4.26*
NM14BMHR2	6.07	3.41*	1.04*	0.31	0.16*	0.93*	0.44*	0.32	3.18	4.23*
AFX148091	5.50	3.43*	0.92	0.45*	0.14*	0.98*	0.50*	0.42*	3.45*	4.13*
57Q53	5.95	2.94	0.93	0.37	0.11*	0.84	0.38	0.33	2.97	3.95
NM14MLLS2	5.83	2.97	0.97*	0.30	0.13*	0.85	0.33	0.22	2.82	3.91
Hi-Gest 660	5.69	2.99	0.83	0.40	0.12*	0.80	0.33	0.26	2.73	3.79
Sandpiper	5.64	2.94	0.90	0.29	0.09*	0.79	0.38	0.30	2.72	3.77
Cisco II	5.72	2.75	0.89	0.23	0.05*	0.78	0.21	0.15	2.26	3.49
WL 440HQ	5.32	2.71	0.78	0.21	0.04*	0.73	0.31	0.22	2.32	3.43
Mean	6.32	3.38	0.96	0.38	0.14	0.92	0.44	0.36	3.20	4.30
LSD (0.05)	1.29	0.65	0.15	0.16	NS	0.21	0.26	0.19	0.95	0.79
CV%	13.93	13.54	10.77	29.40	77.18	15.83	41.87	36.87	21.07	22.47

†Data were analyzed using analysis of covariance where check plots of NuMex Bill Melton were used as the covariate.

2015 Harvest dates: 30-Apr, 6-Jun, 7-Jul, 13-Aug, 10-Sep, and 28-Oct.

2016 Harvest dates: 27-Apr, 27-May, 30-Jun, 27-Jul, 13-Sep, and 21-Oct.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 4. Dry matter yields (tons/acre) of alfalfa varieties sown October 7, 2014, at NMSU's Leyendecker Plant Science Research Center at Las Cruces and flood-irrigated twice per cutting from March 7 until June 13 to apply 39.42 inches, after which irrigation was terminated (irrigation termination)†.

Variety Name	2015 Total	2016 Total	2017 Harvests			2017 Total	3-Yr Average
			26-Apr	25-May	27-Jun		
NM14BMHS1	7.15**	3.76**	1.27*	0.72	0.77*	2.77*	4.55**
NM14BMHR2	6.96*	3.73*	1.33*	0.73*	0.73*	2.78*	4.49*
AmeriStand 803T	6.63*	3.61*	1.20*	0.84*	0.99*	3.04*	4.43*
NM14BMC0	7.08*	3.59*	1.21*	0.65	0.72*	2.58*	4.42*
NM14MalHS3	6.89*	3.60*	1.28*	0.70	0.74*	2.72*	4.41*
NM1407227	6.69*	3.56*	1.38**	0.75*	0.83*	2.95*	4.40*
NM14BMHS3	6.79*	3.55*	1.23*	0.71	0.81*	2.76*	4.35*
NM14GTAF07235	6.18	3.74*	1.22*	0.86*	1.05*	3.14**	4.34*
NM14BM1008251	6.60	3.64*	1.19*	0.79*	0.79*	2.77*	4.34*
AFX148091	6.45	3.49*	1.12*	0.81*	0.99*	2.92*	4.29*
Cisco II	6.39	3.48*	1.20*	0.73*	0.89*	2.82*	4.24*
NM14MLLS2	6.80*	3.47*	1.13*	0.61	0.63*	2.37*	4.22*
NuMex Bill Melton	6.30	3.51*	1.18*	0.76*	0.85*	2.79*	4.18*
Wilson	6.16	3.50*	1.22*	0.78*	0.85*	2.84*	4.18*
Sandpiper	6.27	3.54*	1.16*	0.72	0.84*	2.72*	4.18*
4N900	6.48	3.42*	1.10*	0.73*	0.77*	2.61*	4.17*
Malone	5.63	3.62*	1.12*	0.88**	1.11**	3.12*	4.13*
NM14ALWLHQ	6.65	3.42*	1.02*	0.63	0.62*	2.26*	4.12*
FSG903	6.61	3.33*	1.05*	0.70	0.67*	2.43*	4.11*
AFX149092	6.47	3.37*	1.00*	0.72	0.68*	2.39*	4.10*
Hi-Gest 660	6.45	3.25*	0.92*	0.63	0.68*	2.24*	3.98*
WL 656HQ	6.23	3.21*	0.94*	0.67	0.67*	2.28*	3.92*
WL 440HQ	6.18	3.17*	1.02*	0.63	0.64*	2.29*	3.88*
57Q53	6.02	3.21*	1.06*	0.67	0.68*	2.41*	3.86*
Mean	6.50	3.49	1.15	0.73	0.79	2.67	4.22
LSD (0.05)	0.49	NS	NS	0.16	NS	NS	NS
CV%	5.66	9.62	17.16	15.19	28.17	19.00	14.17

†Data were analyzed using analysis of covariance where check plots of NuMex Bill Melton were used as the covariate.

2015 Harvest dates: 29-May, 12-Jun, 17-Jul, and 12-Aug.

2016 Harvest dates: 28-Apr, 2-Jun, and 29-Jun.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 5. Dry matter yields (tons/acre) of sprinkler-irrigated alfalfa varieties sown March 31, 2014, at NMSU's Agricultural Science Center at Artesia†.

Variety Name	2014 Total	2015 Total	2016 Total	2017 Harvests					2017 Total	4-Yr Average
				12-May	9-Jun	19-Jul	22-Aug	10-Oct		
Malone	4.28*	7.97*	9.00*	1.80*	2.29*	1.85*	2.91**	2.29**	11.12**	8.09**
NM Common	4.36**	8.13*	8.47*	1.97*	2.48*	1.99*	2.19*	1.94*	10.56*	7.88*
56S82	4.15*	8.37**	8.49*	1.78*	2.51*	2.03**	2.53*	1.60*	10.44*	7.86*
SW 7410	4.18*	7.69*	8.82*	1.97*	2.74**	1.89*	2.48*	1.52*	10.58*	7.82*
SW 8421S	3.82*	8.09*	8.49*	1.92*	2.32*	1.80*	2.82*	1.99*	10.85*	7.81*
Artesian Sunrise	4.16*	7.56*	8.77*	2.44**	2.60*	1.96*	2.36*	1.36*	10.72*	7.80*
NuMex Bill Melton	4.06*	8.03*	8.93*	1.94*	2.27*	1.86*	2.43*	1.65*	10.15*	7.79*
Wilson	4.30*	7.84*	9.10*	1.74*	2.31*	1.91*	2.34*	1.60*	9.90*	7.78*
58N57	4.42*	7.57*	9.41**	1.90*	2.34*	1.92*	2.32*	1.24*	9.71*	7.78*
SW 8357	4.09*	8.00*	8.71*	1.90*	2.66*	1.48*	1.94*	1.52*	9.49*	7.57*
WL 535HQ	4.08*	7.51*	8.79*	1.99*	2.29*	1.71*	2.25*	1.50*	9.74*	7.53*
DG9212	4.45*	7.66*	8.21*	1.90*	2.36*	1.85*	2.19*	1.45*	9.74*	7.51*
SW 8208	4.49*	7.50*	8.24*	2.08*	2.04	1.62*	2.32*	1.64*	9.70*	7.48*
Zia	4.32*	7.37*	8.54*	1.72*	1.95	1.83*	2.36*	1.81*	9.67*	7.47*
Dona Ana	3.98*	7.41*	8.37*	1.90*	2.47*	1.41*	2.41*	1.92*	10.11*	7.47*
African Common	4.28*	7.18*	8.57*	1.81*	2.04	1.61*	2.21*	2.09*	9.76*	7.45*
55VR05	3.68*	7.00*	7.93*	2.12*	1.71	1.47*	2.35*	1.55*	9.20*	6.95*
Mean	4.18	7.70	8.64	1.93	2.31	1.78	2.38	1.69	10.08	7.65
LSD (0.05)	NS	NS	NS	NS	0.53	NS	NS	NS	NS	NS
CV%	9.91	9.50	7.57	19.39	16.09	18.55	24.22	24.71	12.00	11.07

†Data were analyzed using analysis of variance.

2014 Harvest dates: 24-Jun, 2-Sep, and 3-Oct.

2015 Harvest dates: 26-May, 13-Jul, 12-Aug, and 30-Sep.

2016 Harvest dates: 8-Jun, 11-Jul, 8-Aug, 30-Sep, and 28-Nov.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 6. Dry matter yields (tons/acre) of sprinkler-irrigated alfalfa varieties sown September 16, 2016, at NMSU's Agricultural Science Center at Artesia†.

Variety Name	2017 Harvests					2017 Total
	16-May	22-Jun	21-Jul	24-Aug	16-Oct	
SW 7408	2.10**	2.23*	1.83*	1.91**	1.31**	9.41**
NuMex Bill Melton	2.02*	2.32**	1.76*	1.79*	1.30*	9.16*
MS sunstra 155203	1.72*	2.08*	2.01**	1.82*	1.29*	8.67*
SW 8476	1.75*	2.15*	1.60	1.78*	1.16*	8.29*
SW 8412	1.75*	1.75	1.68	1.74*	1.28*	8.09*
SW 8409	1.54	2.02*	1.65	1.54*	1.23*	8.08*
African Common	1.69	1.93*	1.51	1.60*	1.12*	8.05*
MS sunstra 155204	1.72*	2.09*	1.22	1.69*	1.26*	7.99*
Dona Ana	1.74*	1.84	1.49	1.63*	1.09*	7.82*
55VR08	1.41	2.06*	1.45	1.74*	1.23*	7.78*
SW 7473	1.67	1.71	1.39	1.73*	1.19*	7.78*
Zia	1.50	1.96*	1.34	1.74*	0.99*	7.64*
New Mexico 11-1	1.49	1.67	1.44	1.65*	1.21*	7.63*
NM Common	1.48	1.65	1.52	1.63*	1.18*	7.36*
Mean	1.68	1.96	1.56	1.71	1.20	8.13
LSD (0.05)	0.39	0.41	0.28	NS	NS	NS
CV%	15.99	14.75	12.40	14.15	11.57	10.50

†Data were analyzed using analysis of covariance where check plots of AmeriStand 803T were used as the covariate.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 7. Dry matter yields (tons/acre) of alfalfa varieties sown May 12, 2015, at NMSU's Agricultural Science Center at Tucumcari and sprinkler-irrigated twice per week with treated municipal wastewater†.

Variety Name	2016 Total	2017 Harvests						2017 Total	2-Yr Average
		19-May	20-Jun	19-Jul	15-Aug	11-Sep	30-Oct		
NuMex Bill Melton	4.62**	1.93*	1.10*	0.36*	1.04*	0.97*	0.46*	5.85**	5.23**
6829R	4.38*	1.72*	1.21**	0.39*	1.06*	0.94*	0.38*	5.68*	5.03*
NM14BMHS1	4.19*	1.75*	1.02*	0.43*	0.90*	0.82	0.46*	5.36*	4.78*
NM14BMHR2	4.17*	1.84*	0.93	0.36*	1.05*	0.75	0.41*	5.33*	4.75*
NM14BMC0	3.99*	1.88*	0.94	0.27*	0.97*	0.74	0.46*	5.26*	4.62*
Mallard	3.98*	1.87*	1.00*	0.26*	1.00*	0.73	0.36	5.23	4.61*
NM14BM1008251	3.94*	1.27*	1.14*	0.44**	0.90*	0.82	0.44*	5.00	4.47
Malone	3.77	1.40*	0.96	0.40*	1.09*	0.93*	0.44*	5.22*	4.49
NM14MalHS3	3.69	1.98**	0.68	0.28*	1.09*	0.69	0.41*	5.12	4.41
African Common	3.65	1.64*	1.12*	0.37*	1.13**	0.83	0.50**	5.59*	4.62
NM14MLLS2	3.65	1.47*	0.84	0.24*	0.88*	0.72	0.35	4.50	4.07
NM Common	3.52	1.60*	0.76	0.31*	1.11*	0.98**	0.35	5.10	4.31
ICON	3.33	1.45*	0.64	0.27*	0.92*	0.79	0.45*	4.51	3.92
SW 5909	3.27	1.56*	0.78	0.24*	0.99*	0.68	0.37	4.61	3.94
SW 5213	3.27	1.63*	0.87	0.28*	1.03*	0.78	0.35	4.93	4.10
Zia	3.23	1.46*	0.93	0.31*	1.06*	0.80	0.38	4.94	4.09
Red Falcon BR	3.19	1.69*	0.65	0.29*	0.85*	0.53	0.19	4.19	3.69
SW 5113	3.10	1.39*	0.75	0.28*	0.99*	0.64	0.25	4.28	3.69
Roadrunner	3.03	1.61*	0.91	0.35*	0.82*	0.54	0.28	4.50	3.76
Mean	3.68	1.64	0.91	0.32	0.99	0.77	0.38	5.01	4.35
LSD (0.05)	0.72	NS	0.24	NS	NS	0.14	0.10	0.72	0.63
CV%	13.85	24.57	18.73	35.41	15.97	12.86	18.96	10.18	14.38

†Data were detrended using nearest neighbor analysis and analyzed using analysis of variance.

2016 Harvest dates: 24-May, 22-Jun, 9-Aug, 13-Sep, and 8-Nov.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 8. Dry matter yields (tons/acre) of flood-irrigated alfalfa varieties sown September 20, 2016, at NMSU's Agricultural Science Center at Los Lunas†.

Variety Name	2017 Harvests				2017 Total
	6-Jun	18-Jul	1-Sep	6-Nov	
NuMex Bill Melton	2.62**	2.16*	1.35*	0.90*	7.03**
WL 454HQ.RR	2.61*	2.15*	1.20*	0.87	6.83*
NM14BMHS1	2.56*	2.00*	1.39*	0.84	6.78*
AmeriStand 715NT RR	2.14*	2.12*	1.56**	0.92*	6.73*
NM14MalHS3	2.52*	2.05*	1.16*	0.97*	6.70*
Artesian Sunrise	2.39*	2.03*	1.17*	1.03*	6.62*
Meadowlark	2.42*	1.90	1.33*	0.93*	6.59*
Roadrunner	2.23*	2.03*	1.34*	0.98*	6.57*
Transition 6.10RR	2.39*	2.01*	1.24*	0.89	6.51*
Archer III	2.06	2.22*	1.26*	0.91*	6.44*
msSunstra 155204	2.16*	2.10*	1.21*	0.95*	6.41*
msSunstra 155206	1.96	2.21*	1.49*	0.75	6.41*
HybriForce 2600	1.96	2.29**	1.28*	0.88	6.41*
Stratica RR	2.23*	1.85	1.37*	0.88	6.32*
SW 7473	2.10*	2.13*	1.36*	0.74	6.32*
WL 440HQ	2.10*	2.07*	1.29*	0.83	6.29*
Dona Ana	2.11*	1.90	1.33*	0.86	6.20
NM14ALWLHQ	2.20*	1.80	1.36*	0.81	6.17
WL 372HQ.RR	1.86	2.13*	1.32*	0.80	6.10
Six Shooter RR	1.95	2.06*	1.29*	0.78	6.08
Hi-Gest 660	2.05	1.83	1.30*	0.86	6.04
Malone	1.83	1.93	1.39*	0.83	5.97
Tonnica RR	2.10*	1.89	1.19*	0.78	5.95
SW 7408	2.00	1.82	1.28*	0.85	5.95
WL 552HQ.RR	1.58	2.04*	1.16*	1.04**	5.83
AmeriStand 855T RR	1.91	1.85	1.20*	0.79	5.75
NM Common	1.89	1.80	1.14*	0.76	5.59
NM14BM1008251	1.76	1.77	1.25*	0.77	5.54
Mean	2.13	2.00	1.29	0.86	6.29
LSD (0.05)	0.55	0.31	NS	0.15	0.76
CV%	18.44	11.02	14.35	12.05	8.61

†Data were detrended using nearest neighbor analysis and analyzed using analysis of variance.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 9. Dry matter yields (tons/acre) of sprinkler-irrigated alfalfa varieties sown August 21, 2014, at NMSU's Agricultural Science Center at Farmington†.

Variety Name	2015 Total	2016 Total	2017 Harvests			2017 Total	3-Yr Average
			7-Jun	11-Jul	24-Aug		
Raven	10.48*	10.13*	2.25*	2.69*	2.35**	7.29*	9.30**
Ranger	10.23*	10.31*	2.24*	2.77**	2.26*	7.26*	9.27*
MagnaGraze II	9.87*	10.50**	2.50**	2.55*	2.32*	7.36**	9.24*
Arrowhead II	10.18*	10.15*	2.28*	2.61*	2.19*	7.08*	9.14*
Mallard 5	10.50*	10.07*	2.25*	2.58*	1.96*	6.79*	9.12*
Lahontan	10.32*	9.86*	2.31*	2.54*	2.27*	7.11*	9.09*
Roadrunner	10.77**	9.30	1.90	2.30	2.27*	6.47*	8.85*
4S417	10.04*	9.92*	2.12*	2.32*	2.05*	6.49*	8.82*
GrandStand	9.52	9.90*	2.36*	2.37*	2.07*	6.80*	8.74*
Archer III	10.48*	9.34	1.74	2.36*	1.90*	5.99	8.60*
Mountaineer 2.0	9.81*	9.61*	2.24*	2.13	1.94*	6.31	8.58*
PGI 424	10.04*	9.37	2.01	2.30	2.01*	6.32	8.58*
WL 363HQ	9.90*	9.98*	1.76	2.08	1.97*	5.81	8.56
NM Common	9.90*	9.35	1.88	2.14	2.16*	6.18	8.47
Hi-Gest 360	9.87*	9.54*	1.83	2.31	1.87*	6.01	8.47
Gunner	9.61	9.18	2.09*	2.28	2.15*	6.52*	8.44
WL 354HQ	9.86*	9.19	1.63	1.96	2.04*	5.62	8.22
54VR03	9.64*	8.75	1.88	2.34*	1.98*	6.20	8.20
Dona Ana	9.52	9.22	2.03	1.78	1.95*	5.76	8.17
Wilson	9.33	8.82	1.57	2.05	2.02*	5.64	7.93
Malone	8.87	8.71	2.13*	1.90	1.99*	6.03	7.87
Zia	8.87	8.12	1.78	1.58	2.07*	5.43	7.47
Mean	9.89	9.51	2.04	2.27	2.08	6.38	8.60
LSD (0.05)	1.06	0.98	0.46	0.46	NS	0.96	0.73
CV%	7.55	7.26	16.00	14.20	15.00	10.67	10.43

†Data were detrended using nearest neighbor analysis, and analyzed using analysis of variance.

2015 Harvest dates: 2-Jun, 6-Jul, 17-Aug, and 9-Oct.

2016 Harvest dates: 2-Jun, 11-Jul, 17-Aug, and 5-Oct.

**Highest numerical value in the column.

*Not significantly different from the highest numerical value in the column based on the 5% LSD.

NS means that there were no significant differences between the varieties within that column at the 5% level.

Table 10. Characteristics and performance of alfalfa varieties across years and tests in New Mexico.		Varietal Characteristics ¹												Las Cruces									Artesia				Tucumcari		Los Lunas		Farmington					
														2014 ²									2014			2016		2015		2016	2014					
		Pest resistance												N ³			D			ET			2014				2016		2015		2016	2014				
		Variety	Proprietor	RR	FD	WS	Or	BW	FW	AN	PRR	SAA	PA	BAA	SN	RKN	15 ⁴	16	17	15	16	17	15	16	17	14	15	16	17	17	16	17	17	15	16	17
Arrowhead II	Alforex Seeds		2	2		HR	HR	HR	HR	n/r	R	n/r	HR	n/r																				*	*	*
Hi-Gest 360	Alforex Seeds		3	n/r		HR	HR	HR	HR	n/r	n/r	R	n/r	n/r																			*	*	*	
MagnaGraze II	Sharp Brothers		3	2		HR	HR	HR	n/r	R	n/r	n/r	R	n/r																		*	**	*		
Ranger	USDA/Univ. of Nebraska		3	n/r		R	n/r	n/r	n/r	R	n/r	n/r	R	n/r																		*	*	*		
4S417	Mycogen Seeds		4	n/r		n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r																		*	*	*		
54VR03	Pioneer HiBred Int'l	Y	4	n/r		HR	HR	HR	HR	n/r	HR	n/r	n/r	n/r																		*				
GrandStand	Crop Production Services		4	2		HR	HR	HR	HR	R	HR	n/r	MR	n/r																		*	*	*		
Meadowlark	Blue River Hybrids		4	2	Y	HR	HR	HR	HR	R	HR	n/r	R	n/r																	*					
PGI 424	Alforex Seeds		4	2		HR	HR	HR	HR	R	R	n/r	R	n/r																		*		*		
Raven	Blue River Hybrids		4	2	Y	HR	R	R	R	R	HR	R	R	n/r																	*	*	**			
Red Falcon BR	Blue River Hybrids		4	2	Y	HR	HR	HR	HR	n/r	n/r	n/r	R	n/r																						
Roadrunner	Blue River Hybrids		4	2	Y	HR	HR	HR	HR	HR	LR	n/r	R	n/r																	*	**	*	*		
RR Stratica	Croplan Genetics	Y	4	n/r		HR	HR	HR	HR	HR	R	n/r	R	n/r																	*					
SW 4113	S & W Seeds		4	n/r		n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r									*	*												
WL 354HQ	W-L Research		4	1		HR	HR	HR	HR	HR	HR	n/r	R	n/r																		*				
55VR05	Pioneer HiBred Int'l	Y	5	n/r		HR	HR	HR	HR	n/r	n/r	n/r	HR	n/r									*	*	*	*										
55VR08	Pioneer HiBred Int'l	Y	5	2		R	HR	HR	HR	R	HR	n/r	n/r	n/r														*								
Archer III	America's Alfalfa		5	2		HR	HR	HR	HR	n/r	HR	n/r	HR	HR																	*	*		*		
Gunner	Croplan Genetics		5	1		HR	HR	HR	HR	HR	R	n/r	R	n/r																						
Mallard 5	Blue River Hybrids		5	2	Y	HR	HR	HR	HR	R	HR	n/r	R	n/r															*			*	*	*		
Mountaineer 2.0	Croplan Genetics		5	2		HR	HR	HR	HR	R	HR	n/r	HR	R																	*	*	*	*		
RR Tonnica	Croplan Genetics	Y	5	n/r		HR	HR	HR	HR	n/r	R	n/r	R	n/r																						
SW 5213	S & W Seeds		5	n/r		n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r									*	*												
SW 5909	S & W Seeds		5	n/r		n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r									*	*												
WL 363HQ	W-L Research		5	2		HR	HR	HR	HR	R	R	MR	MR	MR																	*	*				
WL 372HQ.RR	W-L Research	Y	5	2		HR	HR	HR	HR	HR	HR	n/r	HR	n/r																						

¹RR=Roundup Ready if "Y"; FD=Fall Dormancy (1 & 2 Very dormant; 3 & 4 Dormant; 5 Moderately dormant; 6 & 7 Semi-dormant; 8 & 9 Non-dormant; 10 & 11 Very non-dormant), WS=Winter Survival (1=No injury, 6=Dead plants),

Or=Organically certified seed, BW=Bacterial wilt, PRR=Phytophthora root rot, FW=Fusarium wilt, AN=Anthracnose, SAA=Spotted alfalfa aphid, PA=Pea aphid, BAA=Blue alfalfa aphid, SN=Stem nematode, RKN=Rootknot nematode

(southern); (S=Susceptible, LR=Low resistance, MR=Moderate resistance, R=Resistant, HR=High resistance, n/r indicates either that the variety was not rated for that characteristic or no rating was available).

²Establishment year.

³N, D, and ET signify normal irrigation, drought irrigation, and early termination irrigation, respectively.

⁴Harvest year.

Shaded boxes indicate that the variety was not in the test.

***Significantly higher than all other values in the column.

**Highest yielding variety in the test for that year, except for Los Lunas 2013 test when it is the second highest yielding variety.

*Not significantly different from the highest yielding variety in the test for that year, except for Los Lunas 2013 test in 2014 when it is not significantly different from the second highest yielding variety.

L.M. Lauriault, I.M. Ray, C.A. Pierce, O. Burney, R.P. Flynn, M.A. Marsalis, M.K. O'Neill, A.E. Cunningham, C. Havlik, and M.M. West

New Mexico St. Univ. College of Agricultural, Consumer and Environmental Sciences. Agric. Exp. Stn and Coop. Ext. Ser.

Table 10 (cont.). Characteristics and performance of alfalfa varieties across years and tests in New Mexico.		Varietal Characteristics ¹												Las Cruces									Artesia				Tucumcari		Los Lunas	Farmington							
														2014 ²									2014			2016		2015		2016	2014						
		Pest resistance												N ³			D			ET			2014				2016		2015		2016	2014					
		Variety	Proprietor	RR	FD	WS	Or	BW	FW	AN	PRR	SAA	PA	BAA	SN	RKN	15 ⁴	16	17	15	16	17	15	16	17	14	15	16	17	17	16	17	17	15	16	17	
56S82	Pioneer HiBred Int'l		6	5		HR	HR	HR	HR	HR	HR	HR	HR	HR										*	**	*	*										
Cisco II	Alforex Seeds		6	2		HR	HR	HR	R	n/r	HR	n/r	R	R	*	*						*	*														
Hi-Gest 660	Alforex Seeds		6	n/r		R	HR	HR	R	n/r	n/r	R	n/r	n/r		*						*	*														
HybriForce-2600	Alforex Seeds		6	2		HR	HR	HR	HR	n/r	R	n/r	HR	R																			*				
ICON	S & W Seeds		6	n/r		n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r																							
Lahontan	USDA/Univ. of Nevada		6	n/r		R	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r																			*	*	*		
msSunstra 155203	Alforex Seeds		6	n/r		n/r	n/r	n/r	n/r	n/r	n/r	HR	n/r	n/r													*										
msSunstra 155204	Alforex Seeds		6	n/r		n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r													*						*				
msSunstra 155206	Alforex Seeds		6	n/r		n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r	n/r																			*				
Six Shooter RR	Croplan Genetics	Y	6	n/r		HR	HR	HR	HR	HR	HR	n/r	HR	R																							
Wilson	NMSU/Roswell Seed		6	n/r		R	R	n/r	n/r	MR	R	n/r	MR	n/r		*		*	*	*		*	*	*	*	*	*	*									
WL 440HQ	W-L Research		6	n/r		HR	HR	HR	HR	HR	HR	n/r	HR	HR		*						*	*										*				
WL 454HQ.RR	W-L Research	Y	6	n/r		R	HR	HR	HR	HR	HR	n/r	HR	n/r																			*				
57Q53	Pioneer Hi-Bred Int'l.		7	n/r		MR	HR	HR	R	MR	HR	R	R	MR		*						*	*														
6829R	Nexgrow Alfalfa		7	n/r		R	R	HR	HR	HR	HR	R	HR	n/r														*	*								
Ameristand 715NTRR	America's Alfalfa	Y	7	n/r		R	HR	HR	HR	HR	HR	n/r	HR	n/r																			*				
Artesian Sunrise	Croplan Genetics		7	n/r		MR	HR	R	HR	HR	HR	R	R	n/r										*	*	*	*							*			
Dona Ana	NMSU/Roswell Seed		7	n/r		MR	MR	LR	R	MR	R	n/r	n/r	n/r										*	*	*	*	*									
Malone	NMSU/Roswell Seed		7	n/r	Y	R	HR	R	R	R	HR	S	MR	n/r		*	*	*	*	*	*	*	*	*	*	*	*	*			*						
NuMex Bill Melton	New Mexico State University		7	n/r		MR	R	R	R	R	MR	MR	n/r	n/r		*	**	*	*	**		*	*	*	*	*	*	*	*	**	**	**	**	**	**	**	**
SW 7408	S & W Seeds		7	n/r		R	n/r	HR	HR	n/r	n/r	n/r	R	n/r														**									
SW 7410	S & W Seeds		7	n/r		R	R	HR	MR	HR	R	R	MR	R										*	*	*	*										
SW 7473	S & W Seeds		7	n/r		R	n/r	HR	HR	n/r	n/r	n/r	HR	n/r													*						*				
Transition 6.10RR	Croplan Genetics	Y	7	n/r		R	R	R	HR	R	HR	n/r	MR	n/r														*					*				
58N57	Pioneer HiBred Int'l		8	n/r		LR	R	HR	HR	R	HR	HR	MR	HR										*	*	**	*										
AmeriStand 803T	America's Alfalfa		8	n/r		MR	HR	m	h	r	HR	HR	HR	HR	**	**	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*

¹RR=Roundup Ready if "Y"; FD=Fall Dormancy (1 & 2 Very dormant; 3 & 4 Dormant; 5 Moderately dormant; 6 & 7 Semi-dormant; 8 & 9 Non-dormant; 10 & 11 Very non-dormant), WS=Winter Survival (1=No injury, 6=Dead plants), Or=Organically certified seed, BW=Bacterial wilt, PRR=Phytophthora root rot, FW=Fusarium wilt, AN=Anthracnose, SAA=Spotted alfalfa aphid, PA=Pea aphid, BAA=Blue alfalfa aphid, SN=Stem nematode, RKN=Rootknot nematode (southern); (S=Susceptible, LR=Low resistance, MR=Moderate resistance, R=Resistant, HR=High resistance, n/r indicates either that the variety was not rated for that characteristic or no rating was available).

²Establishment year.

³N, D, and ET signifiyormal irrigation, drought irrigation, and early termination irrigation, respectively.

⁴Harvest year.

Shaded boxes indicate that the variety was not in the test.

***Significantly higher than all other values in the column.

**Highest yielding variety in the test for that year, except for Los Lunas 2013 test when it is the second highest yielding variety.

*Not significantly different from the highest yielding variety in the test for that year, except for Los Lunas 2013 test in 2014 when it is not significantly different from the second highest yielding variety.

L.M. Lauriault, I.M. Ray, C.A. Pierce, O. Burney, R.P. Flynn, M.A. Marsalis, M.K. O'Neill, A.E. Cunningham, C. Havlik, and M.M. West
 New Mexico St. Univ. College of Agricultural, Consumer and Environmental Sciences. Agric. Exp. Stn and Coop. Ext. Ser.

Table 11. New Mexico State University Agricultural Experiment Station and Cooperative Extension Service publications related to alfalfa management.

Number	Title
A-123	Sampling for plant tissue analysis
A-129	Nitrogen fixation by legumes
A-130	Inoculation of legumes
A-131	Certified seed
A-137	Soil analysis: A key to soil nutrient management
A-145	Certified noxious weed free program
A-229	Phymatotrichum root rot
A-325	Managing weeds in alfalfa
A-326	Downy mildew on alfalfa
A-333	User manual of the alfalfa yield predictor
A-334	Beet armyworm in New Mexico Hay
A-335	Variegated cutworm in New Mexico Hay
A-336	Managing Roundup Ready alfalfa and conventional or organic alfalfa hay in nearby fields in New Mexico
A-337	Recommendations for Roundup Ready alfalfa weed management and stand removal in New Mexico
A-338	Alfalfa weevil control options in New Mexico
A-339	Alfalfa integrated pest management: Aphids
H-158	How to collect and send plant specimens for disease diagnosis
CR-536	Blister beetles in alfalfa
CR-633	Using a computer application to predict irrigated alfalfa yield
CR-641	Hay quality, sampling and testing
CR-644	Assessing alfalfa stands after winter injury, freeze damage, or any time renovation is considered in New Mexico
CR-646	Managing alfalfa during drought
CR-654	Selecting alfalfa varieties for New Mexico
CR-659	Whitefringed beetle in New Mexico alfalfa
CR-668	Reducing harvest and post-harvest losses of alfalfa and other hay
RR-766	Furrow-irrigated alfalfa dry matter yield is not affected by different seeding rates in the Southern High Plains, USA
RR-772	Observations on how cowpea aphid affects alfalfa

These publications, and alfalfa variety test reports from previous years, are available from your county office of the NMSU Cooperative Extension Service or online at <http://forages.nmsu.edu/resources.html> and aces.nmsu.edu/pubs/



New Mexico State University
All About Discovery!TM

The College of Agricultural, Consumer and Environmental Sciences is an engine for economic and community development in New Mexico, improving the lives of New Mexicans through academic, research, and extension programs.

New Mexico State University is an equal opportunity/affirmative action employer and educator. NMSU and the U.S. Department of Agriculture cooperating.